

NMR Applications in Formation Evaluation: Well Logging and Core Characterization Excerpted from the "Bibliography of Well Logging Applications"

Updated through 12/97 compiled by Stephen Prensky January 30, 1998
U.S. Minerals Management Service, New Orleans, Louisiana

(For additional references published prior to 1990 refer to Jackson and Mathews, 1993; Gladden, 1994)

1. Abdel-Latif, A., Sungur, D., Stockden, I., Cao Minh, C., 1997, Borehole nuclear magnetic resonance experience and reservoir applications in west Kuwait carbonate reservoirs, SPE-37771, in SPE Middle East oil show and exhibition proceedings: Society of Petroleum Engineers. Also published in 1997 in condensed form: Journal of Petroleum Technology, v. 49, no. 11, p. 1247-1249.
2. Abragam, A., 1961, The principles of nuclear magnetism: Oxford University Press.
3. Aguilera, R., 1988, A new approach for analysis of the nuclear magnetic log-resistivity combination, paper 88-39-109, in 39th annual technical meeting, preprints, volume 3: Petroleum Society of CIM [Canadian Institute of Mining], Calgary, p. 12 p. Later published in 1990: Journal of Canadian Petroleum Technology, v. 29, no. 1, January-February, p. 67-71.
4. Ahmed, U., Crary, S.F., and Coates, G.R., 1989, Permeability estimation; the various sources and their interrelationship, SPE-19604, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 649-662. Later published in 1990 as, The interrelationship between various permeability measurements, paper I, in International well logging symposium transactions: Society of Professional Well Log Analysts, Beijing Chapter, 21 p. Later published in 1991: Journal of Petroleum Technology, v. 43, no. 5, p. 578-587. Later reprinted in 1994, in Petrophysics: Society of Petroleum Engineers Reprint Series No. 39, p. 118-127.
5. Akbar, M., Petricola, M., Watfa, M., Badri, M., Charara, M., Boyd, A., Cassell, B., Nurmi, R., Delhomme, J.-P., Grace, M., Kenyon, B., and Roestenburg, J., 1995, Classic interpretation problems--evaluating carbonates: Schlumberger Oilfield Review, v. 7, no. 1, p. 4-22
6. Akkurt R., 1990, Effects of motion in pulsed NMR logging: Colorado School of Mines, Golden, unpublished Ph.D. dissertation, 165 p.
7. Akkurt, R., Guillory, A.J., Vinegar, H.J. and Tutunjian, P.N., 1995, NMR logging of natural gas reservoirs, paper N, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p. Later published in 1996, as Akkurt, R., Vinegar, H.J., Tutunjian, P.N., and Guillory, A.J.: The Log Analyst, v. 37, no. 5, p. 33-42.
8. Akkurt, R., Moore, A., and Freeman, J., 1997, Impact of NMR in the development of a deepwater turbidite field, paper SS, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 14 p.
9. Akkurt, R., Prammer, M., and Moore, A., 1996, Selection of optimal acquisition parameters for MRIL logs, paper TT, in 37th annual logging symposium transactions: Society of Professional Well Log Analysts, 13 p. Later published in 1996: The Log Analyst, v. 37, no. 6, p. 43-52.
10. Akkurt, R., Vinegar, H.J. Tutunjian, P.N., and Guillory, A.J., 1995, NMR logging of natural gas reservoirs, paper N, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p.

11. Akselrod, S.M., Danevich, V.I., Saikhov, D.M., and Mamed-zadeh, V.A., 1996, Downhole lateral nuclear magnetic sounding: *The Log Analyst*, v. 37, no. 4, p. 25-29
12. Akselrod, S.M., and Neretin, V.D., 1990, Application of nuclear magnetic resonance in petroleum geology and geophysics [in Russian]: Nedra, Moscow, 191 p.
13. Allen, D., Coates, G., and others, 1988, Probing for permeability--An introduction of measurements: *The Technical Review*, v. 36, no. 1, p. 6-20.
14. Allen, D., Crary, S., Freedman, B., Andreani, M., Klopff, W., Badry, R., Flaum, C., Kenyon, B., Kleinberg, B., Gossenberg, P., Horkowitz, J., Logan, D., Singer J., and White, J., 1997, How to use borehole nuclear magnetic resonance: *Schlumberger Oilfield Review*, v., 9, no. 2, p. 2-15.
15. Altunbay, M., Edwards, C.M., and Georgi, D.T., 1997, Reducing uncertainty in reservoir description with NMR-log data, SPE-38027, in Asia Pacific oil and gas conference proceedings: Society of Petroleum Engineers, p. 99-108.
16. Altunbay, M., Georgi, D., and Takezaki, H., 1997, Improving Swr and permeability estimates in carbonates with hydraulic units and NMR-variable T2-partitioning coefficients, paper I, in 3rd well logging symposium of Japan proceedings: Society of Professional Well Log Analysts, Japan Chapter, 9 p.
17. Artus, D.S., 1967, Nuclear magnetism logging in the Rocky Mountain area: *The Log Analyst*, v. 8, no. 2, p. 16-22.
18. Artus, D.S., 1969, Nuclear magnetism log can find porosity, water-cut: *World Oil*, v. 169, no. 7, p. 89-90, 92.
19. Attard, J.J., Carpenter, T.A., Hall, L.D., Davies, S., Taylor, M.J., and Packer, K.J., 1991, Spatially resolved T1 relaxation measurements in reservoir cores: *Magnetic Resonance Imaging*, v. 9, p. 815-819.
20. Attard, J.J., Doran, S.J., Herrod, N.J., Carpenter, T.E., and Hall, L.D., 1992, Quantitative NMR spin-lattice-relaxation imaging of brine in sandstone reservoir cores: *Journal of Magnetic Resonance*, v. 96, p. 514-525.
21. Attard, J.J., McDonald, P.J., Roberts, S.P., and Taylor, T., 1994, Solid state NMR imaging of irreducible water in reservoir cores for spatially resolved pore surface relaxation estimation: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 355-359.
22. Austin, J.M., and Faulkner, T.L., 1993, Magnetic resonance imaging log evaluates low-resistivity pay: *American Oil and Gas Reporter*, v. 36, no. 8, p. 52-57.
23. Baldwin, B.A., 1993, Immobile water determination in shaly sandstone, paper SCA-9314, in Annual technical conference proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 14 p. Later published in 1994: *The Log Analyst*, v. 35, no. 3, p. 31-37.
24. Baldwin, B.A., and Yamanashi, W.S., 1986, Detecting fluid movement and isolation in reservoir cores using medical NMR imaging techniques, SPE/DOE-14884, in ?th SPE/DOE enhanced oil recovery symposium proceedings, v. 1: Society of Petroleum Engineers, p. 39-48. Later published in 1989: *SPE Reservoir Engineering*, v. 4, no. 2, p. 207-212.
25. Baldwin, B.A., and Yamanashi, W.S., 1988, NMR imaging of fluid dynamics in reservoir core: *Magnetic Resonance Imaging*, v. 6, no. 5, p. 493-500.
26. Baldwin, B.A., and Yamanashi, W.S., 1989, Determining fluid saturation distribution in cores using NMR imaging, SCA-8714, part I, paper N, in Annual technical conference preprints, v. 1 (1987-1989): Society of Professional Well Log Analysts, Society of Core Analysts, Chapter-at-Large, 24 p. Later published in 1991: *The Log Analyst*, v. 32, no. 5, p. 536-549.
27. Baldwin, B.A., and Yamanashi, W.S., 1991, Capillary-pressure determination from NMR images of centrifuged cores plus--Berea Sandstone: *The Log Analyst*, v. 32, no. 5, p. 550-556.
28. Banavar, J.R., and Schwartz, L.M., 1987, Magnetic resonance as a probe of permeability in porous media: *Physical Review Letters*, v. 58, no. 14, p. 1411-1414.
29. Banavar, J.R., and Schwartz, L.M., 1989, Probing porous media with nuclear magnetic resonance, in Klaffer, J., and Drake, J.M., eds., *Molecular dynamics in restricted geometries*: Wiley, New

- York, p. 273-310.
30. Basan, P., Howard, J., Lahann, R., Prichard, T., Attard, J., and Camden, D., 1994, Variations in NMR relaxivity and its relationship to pore size in sandstones from the South Morecambe Field, Irish Sea, paper AA, in 16th European formation evaluation symposium transactions: Society of Professional Well Log Analysts, Aberdeen Chapter, 39 p.
 31. Basan, P. Lowden, B., and Stromberg, S., 1997, Developing strategies using core analysis for calibrating the NMR log: DiaLog [London Petrophysical Society Newsletter], v. 5, no. 6, p. 4-6.
 32. Bendel, P., 1990, Spin-echo attenuation by diffusion in nonuniform field gradients: Journal of Magnetic Resonance, v. 86, p. 509-515.
 33. Bergman, D.J., and Dunn, K.J., 1994, Theory of diffusion in a porous medium with applications to pulsed-field-gradient NMR: Physical Review, Series B, v. 50, p. 9153.
 34. Bergman, D.J., and Dunn, K.J., 1995, NMR of diffusing atoms in a periodic porous medium in the presence of a nonuniform magnetic field: Physical Review, Series E, v. 52, no. 6, p. 6516.
 35. Bergman, D.J., Dunn, K.J., and LaTorraca, G.A., 1995, Magnetic susceptibility contrast and fixed field gradient effects on the spin-echo amplitude in a periodic porous medium diffusion: Bulletin of the American Physical Society, v. 40, p. 695.
 36. Bergman, D.J., Dunn, K.-J., and LaTorraca, G.A., 1996, Magnetic field nonuniformities and NMR of protons diffusing in a porous medium: Magnetic Resonance Imaging, v. 14, no. 7/8, p. 851-855.
 37. Best, D.L., Couturie, S.H., and Herrick, R.C., 1979, An improved nuclear magnetism logging system and its application to formation evaluation, SPE-8361: Society of Petroleum Engineers, 54th Annual Technical Conference Preprint, 8 p.
 38. Bilardo, U., Borgia, G.C., Bortolotti, V., Fantazzini, P., and Mesini, E., 1991, Magnetic resonance lifetimes as a bridge between transport and structural properties of natural porous media: Journal of Petroleum Science and Engineering, v. 5, p. 273-283.
 39. Blackband, S., Mansfield, P., Barnes, J.R., Clague, A.D.H., and Rice, S.A., 1984, Discrimination of crude oil and water in sand and in bore cores using NMR imaging, SPE-13401: Society of Petroleum Engineers, unsolicited manuscript, 17 p. Later published in 1986: SPE Formation Evaluation, v. 1, no. 1, p. 31-34.
 40. Bobroff, S., Guillot, G., Riviere, C., Cuiec, L., and Roussel, J.C., 1996, Quantitative porosity profiles and wettability contrast visualization in sandstone by CPMG imaging: Magnetic Resonance Imaging, v. 14, no. 7/8, p. 907-909.
 41. Borgia, G.C., 1994, A new un-free fluid index in sandstones through NMR studies, SPE-28366, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 37-44. Later published in 1996: SPE Formation Evaluation, v. 11, no. 2, p. 89-93.
 42. Borgia, G.C., Bortolotti, V., Brancolini, A., Brown, R.J.S., and Fantazzini, P., 1996, Developments in core analysis by NMR measurements: Magnetic Resonance Imaging, v. 14, no. 7/8, p. 751-760.
 43. Borgia, G.C., Bortolotti, V., Brown, R.J.S., and Fantazzini, P., 1996, A robust method for calculating geometric mean times from multiexponential relaxation data, using only a few data points and only a few elementary operations: Magnetic Resonance Imaging, v. 14, no. 7/8, p. 895-897.
 44. Borgia, G.C., Bortolotti, V., Dattilo, P., and Fantazzini, P., 1997, Quantitative determination of the degree of heterogeneity within an outcropping reservoir by magnetic resonance imaging, SPE-38751, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology, part 2: Society of Petroleum Engineers, p. 485-490.
 45. Borgia, G., Brancolini, A., Brown, R.J.S., Fantazzini, P., and Ragazzini, G., 1994, Water-air saturation changes in restricted geometries studied by proton relaxation: Magnetic Resonance Imaging, v. 12, p. 191-195.
 46. Borgia, G.C., Brancolini, A., Camazi, A., and Maddinelli, G., 1994, Capillary water determination

- in core plugs; a combined study based on imaging techniques and relaxation analysis: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 221-224.
47. Borgia, G.C., Brighenti, G., Fantazzini, P., Fanti, G.D., and Mesini, E., 1990, Specific surface and fluid transport in sandstones through NMR studies, SPE-20921, in European petroleum conference [Europec 90] proceedings: Society of Petroleum Engineers. Later published in 1992: *SPE Formation Evaluation*, v. 7, no. 3, p. 206-210.
 48. Borgia, G.C., Brown, R.J.S., Fantazzini, P., and others, editors, 1991, Proceedings of the first international conference on recent advances in NMR applications to porous media: *Magnetic Resonance Imaging*, v. 9, no. 5, p. 639-888.
 49. Borgia, G.C., Brown, R.J.S., and Fantazzini, P., 1995, The effect of diffusion and susceptibility differences on T2 measurements for fluids in porous media and biological tissues: *Magnetic Resonance Imaging*, v. 14, no. 7/8, p. 731-736.
 50. Borgia, G.C., Brown, R.J.S., and Fantazzini, P., 1995, Scaling of spin-echo amplitudes with frequency, diffusion coefficient, pore size and susceptibility difference for the NMR of fluids in porous media and biological tissues: *Physical Review, Series E*, v. 51, p. 2104.
 51. Borgia, G.C., Brown, R.J.S., and Fantazzini, P., 1996, The effect of diffusion and susceptibility differences on T2 measurements for fluids in porous media and biological tissues: *Magnetic Resonance Imaging*, v. 14, p. 731.
 52. Borgia, G.C., Brown, R.J.S., and Fantazzini, P., 1996, Nuclear magnetic resonance relaxivity and surface-to-volume ratio in porous media with a wide distribution of pore sizes: *Journal of Applied Physics*, v. 79, p. 3656.
 53. Borgia, G.C., Fantazzini, P., Bore, J.C., Smith, M.E., and Strange, J.H., editors, 1994, Proceedings of the second international meeting on recent advances in MR applications to porous media: *Magnetic Resonance Imaging*, v. 12, no. 2.
 54. Borgia, G.C., Fantazzini, P., Fanti, G., and Mesini, E., 1992, Magnetic resonance relaxation study of preferential wettability effects on displacement efficiency in chalk samples: *Journal of Petroleum Science and Engineering*, v. 8, p. 153-159.
 55. Borgia, G.C., Fantazzini, P., Fanti, G., Mesini, E., Terzi, L., and Valdre, G., 1991, A proton relaxation study of immiscible liquid arrangement in microporous structures: *Magnetic Resonance Imaging*, v. 9, no. 5, p. 695-702.
 56. Borgia, G.C., Fantazzini, P., and Mesini, E., 1990, ¹H spin-lattice relaxation as a fingerprint of porous media: *Magnetic Resonance Imaging*, v. 8, p. 435-447.
 57. Borgia, G.C., Fantazzini, P., and Mesini, E., 1991, Wettability effects on oil-water-configurations in porous media--a nuclear magnetic resonance relaxation study: *Journal of Applied Physics*, v. 70, p. 7623-7625.
 58. Bottomley, P.A., 1982, NMR imaging techniques and applications--a review: *Reviews of Scientific Instruments*, v. 53, p. 1319.
 59. Bowers, M.C., 1993, The use of nuclear magnetic resonance, permeability, and diffusion to characterize the porous microstructure of sandstones: University of South Carolina, unpublished Ph.D. thesis, 166 p.
 60. Bowers, M.C., Ehrlich, R., and Clark, R.A., 1994, Determination of petrographic factors controlling permeability using petrographic, image analysis and core data, Satun field, Pattani Basin, Gulf of Thailand: *Marine and Petroleum Geology*, v. 11, no. 2, p. 148-156.
 61. Bowers, M.C., Ehrlich, R., Howard, J.J., and Kenyon, W.E., 1993, Determination of porosity types from NMR data and their relationship to porosity types derived from thin section, SPE-26307: Society of Petroleum Engineers, unsolicited paper, 45 p. Later published in 1995: *Journal of Petroleum Science and Engineering*, v. 13, p. 1-14.
 62. Bowers, M.C., Purkale, J.D., and Gregory, M.D., 1996, Reconciliation of differences in miscible hydrocarbon gas flood using NMR, SPE/DOE-35389, in 10th SPE/DOE symposium on improved oil recovery proceedings, v. 1: Society of Petroleum Engineers, p. 457-468.
 63. Boyd, A., Darling, H., Tabanou, J., Davis, B., Lyon, B., Flaum, C., Klein, J., Sneider, R.M.,

- Sibbit, A., and Singer, J., 1995, The lowdown on low-resistivity pay: Schlumberger Oilfield Review, v. 7, no. 3, p. 4-18.
64. Boyd, A., Head, E., and Gubelin, W.G., 1995, Lithology-independent, high-resolution porosity logs SPE-30561, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 317-324..
 65. Brancolini, A., Cominelli, A., Kulkarni, R.K., and Watson, A., 1997, Spatial distribution of petrophysical parameters on a core scale using magnetic resonance imaging, paper AAA, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 11 p.
 66. Brancolini, A., Gossenberg, P., Lyne, A.M., Mackenzie, I.S., and Radaelli, F., 1995, The use of NMR core analysis in the interpretation of downhole NMR logs, SPE-30559, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 291-300.
 67. Briskin, M., Robbins, J., Riedel, W.R., and Booker, R., 1985, Magnetic resonance imaging analysis of varved marine sedimentary records of the Gulf of California: Geophysical Research Letters, v. 13, no. 8, p. 753-756.
 68. Brown, J.A., Brown, L.F., and Jackson, J.A., 1981, NMR (nuclear magnetic resonance) measurements on western gas sands core, SPE/DOE-9861, in SPE/DOE Low permeability gas reservoirs symposium proceedings: Society of Petroleum Engineers, p. 321-326. Also published in 1981 as Los Alamos National Laboratory Report No. LA-UR-81-736, 16 p.
 69. Brown, J.A., Jackson, J.A., Brown, L.F., Milewski, J.V., and Travis, B.J., 1982, NMR logging tool development--laboratory studies of tight gas sands and artificial porous material: Los Alamos National Laboratory Report No. LA-UR-82-447, 20 p. Also published in 1982 as SPE-10813, in SPE/DOE unconventional gas recovery symposium proceedings: Society of Petroleum Engineers, p. 203-209.
 70. Brown, J.A., Jackson, J.A., and Koelle, A.R., 1985, Western gas sands project Los Alamos NMR well logging tool development: Los Alamos National Laboratory, Report LA-10374-PR, March, 77 p.
 71. Brown, L.D., 1989, NMR imaging--principles and recent progress, SCA-8812, part II, paper L, in Annual technical conference reprints, v. 1 (1987-1989): Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 27 p.
 72. Brown, L.F., and Travis, B.J., 1983, Using diffusion measurements to determine pore-size distributions in porous media: Chemical Engineering Science, v. 38, p. 843.
 73. Brown, R.J.S., 1961, Proton relaxation in crude oils: Nature, v. 189, no. 4762, p. 387-388.
 74. Brown, R.J.S., 1989, Information available and unavailable from multiexponential relaxation data: Journal of Magnetic Resonance, v. 82, p. 539-561.
 75. Brown, R.J.S., 1996, Nuclear magnetism logging at the "Coyote Institute": Magnetic Resonance Imaging, v. 14, no. 7/8, p. 811-817.
 76. Brown, R.J.S., Borgia, G.C., Fantazzini, P., and Mesini, E., 1991, Problems in identifying multimodal distributions of relaxation times for NMR in porous media: Magnetic Resonance Imaging, v. 9, p. 687-693.
 77. Brown, R.J.S., Borgia, G.C., Fantazzini, P., and Mesini, E., 1991, Need for standardization of laboratory measurements in NMR applications to porous media: Magnetic Resonance Imaging, v. 9, no. 5, p. 885-887.
 78. Brown, R.J.S., and Fantazzini, P., 1993, Conditions for initial quasilinear T2-1 vs tau for CPMG NMR with diffusion and susceptibility differences in porous media and tissues: Physical Review, Series B, v. 47, p. 14,823.
 79. Brown, R.J.S., and Fatt, I., 1956, Measurement of fractional wettability of oil fields' rocks by the nuclear magnetic relaxation method, SPE-743-G: Petroleum Branch of AIME [Society of Petroleum Engineers] 31st Annual Fall Meeting Preprints, 4 p..
 80. Brown, R.J.S., and Gamson, B.W., 1960, Nuclear magnetism logging, SPE-1305-G: Journal of Petroleum Technology, v. 12, no. 8, p. 199-207. Also published in 1960: AIME Transactions, v.

- 219, p. 199-207.
81. Brown, R.J.S., and Neuman, C.H., 1980, Processing and display of nuclear magnetism logging signals--Application to residual oil determination, paper K, in 21st annual logging symposium transactions: Society of Professional Well Log Analysts, 25 p.
 82. Brown, R.J.S., and Neuman, C.H., 1982, The nuclear magnetism log--A guide for field use: *The Log Analyst*, v. 23, no. 5, p. 4-9.
 83. Brownstein, K.R., and Tarr, C.E., 1977, Spin-lattice relaxation in a system governed by diffusion: *Journal of Magnetic Resonance*, v. 26, p. 17-24.
 84. Brownstein, K.R., and Tarr, C.E., 1979, Importance of classical diffusion in NMR studies of water in biological cells: *Physical Review, Series A*, v. 19, p. 2446-2453.
 85. Callaghan, P.T., 1991, *Principles of nuclear magnetic resonance microscopy*: Clarendon Press, Oxford, England.
 86. Callaghan, P.T., 1996, NMR imaging, NMR diffraction and applications of pulsed gradient spin echoes in porous media: *Magnetic Resonance Imaging*, v. 14, no. 7/8, p. 701-709.
 87. Callaghan, P.T., and Eccles, C.D., 1987, Sensitivity and resolution in NMR imaging: *Journal of Magnetic Resonance*, v. 71, p. 426-445.
 88. Callaghan, P.T., and Eccles, C.D., 1988, Diffusion-limited resolution in nuclear magnetic resonance microscopy: *Journal of Magnetic Resonance*, v. 78, p. 1-8.
 89. Callaghan, P.T., Coy, A., Halpin, T.P.J., MacGowan, D., Packer, K.J., and Zelaya, F.O., 1992, Diffusion in porous systems and the influence of pore morphology in pulsed gradient spin-echo nuclear magnetic resonance studies: *Journal of Chemical Physics*, v. 97, p. 651-662.
 90. Callaghan, P.T., Coy, A., MacGowan, D., Packer, K.J., and Zelaya, F.O., 1991, Diffraction-like effects in NMR diffusion studies of fluids in porous solids: *Nature*, v. 351, p. 467-469.
 91. Callaghan, P.T., MacGowan, D., Packer, K.J., and Zelaya, F.O., 1991, Influence of field gradient strength in NMR studies of diffusion in porous media: *Magnetic Resonance Imaging*, v. 9, p. 663-671.
 92. Camden, D., 1994, NMR theory made simple: *Dialog [London Petrophysical Society Quarterly Newsletter]*, v. 2, no. 4, October, p. 4-5.
 93. Caprihan, A., and Fukushima, E., 1990, Flow measurements by NMR: *Physical Reports*, v. 198, p. 195-235.
 94. Carati, C., and Maddinelli, G., 1994, NMR spectroscopy and relaxation--powerful tools for estimating residual fluids and petrophysical parameters, paper K, in 16th European formation evaluation symposium transactions: Society of Professional Well Log Analysts, Aberdeen Chapter, 12 p.
 95. Carr, M.B., Ehrlich, R., Bowers, M.C., and Howard, J.J., 1996, Correlation of porosity types derived from NMR data and thin section image analysis in a carbonate reservoir: *Journal of Petroleum Science and Engineering*, v. 14, p. 115-131.
 96. Chandler, R., 1995, NMR hardware history and MRIL, chapter 6, in Georgi, D.T., ed., *Nuclear magnetic resonance logging short course notes*, 36th annual symposium, Paris, France: Society of Professional Well Log Analysts, variously paginated.
 97. Chandler, R.N., Drack, E.D., Miller, M.N., and Prammer, M.G., 1994, Improved log quality with a dual-frequency pulsed NMR tool, SPE-28365, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 23-35.
 98. Chandler, R.N., Kenyon, W.E., Morriss, C.E., 1987, Reliable nuclear magnetism logging--with examples in effective porosity and residual oil saturation, paper C, in 28th annual logging symposium transactions: Society of Professional Well Log Analysts, 25 p.
 99. Chang, C.T., and Edwards, C.M., 1991, Proton NMR two-component chemical shift imaging of fluids in porous media, SCA-9114, in 5th annual technical conference preprints, v. 3: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 15 p. Later published in 1993: *The Log Analyst*, v. 35, no. 6, November-December, p. 20-28.

100. Chang, D., Vinegar, H., Morriss, C., and Straley, C., 1994, Effective porosity, producible fluid and permeability in carbonates from NMR logging, paper A, in 35th annual logging symposium transactions: Society of Professional Well Log Analysts, 21 p. Later published in 1997: *The Log Analyst*, v. 38, no. 2, p. 60-72.
101. Chang, P., Qiao, J., Chen, S., and Watson, T., 1996, Investigating the utility of NMR spectroscopic techniques for fracture characterization, paper QQ, in 37th annual logging symposium transactions: Society of Professional Well Log Analysts, 13 p.
102. Chapman, A., and Knight, R., 1991, Investigating pore geometry with nuclear magnetic resonance, paper, I, in 13th formation evaluation symposium transactions: Canadian Well Logging Society, 22 p.
103. Charдаire, C., and Roussel, J.-C., 1990, NMR imaging of fluid saturation distributions in core samples using a high magnetic field, in Worthington, P.F., ed., *Advances in core evaluation, accuracy, and precision in reserves estimation* [Reviewed proceedings of the first Society of Core Analysts European core analysis symposium, May 21-23, London, UK]: Gordon and Breach Science Publishers, New York, p. 301-315.
104. Charдаire-Riviere, C., Forbes, P., Zhang, J.F., Chavent, G., and Lenormand, R., 1992, Improving the centrifuge technique by measuring local saturations, SPE-24882, in SPE annual technical conference and exhibition proceedings, v. sigma, *Reservoir engineering*: Society of Petroleum Engineers, p. 193-208.
105. Charдаire-Riviere, C., and Roussel, J.-C., 1991, Use of a high magnetic field to visualize fluids in porous media by MRI, SCA-9112, in 5th annual technical conference preprints: Society of Core Analysts, Society of Professional Well Log Analysts Chapter-at-Large, v. 3, 11 p. Also published in 1991: *Magnetic Resonance Imaging*, v. 9, no. 5, p. 827-832.
106. Charдаire-Riviere, C., and Roussel, J.C., 1992, Principle and potential of nuclear magnetic resonance applied to the study of fluids in porous media: *Revue de l'Institut Francais du Petrole*, v. 47, no. 4, p. 503-523.
107. Chen, J., Dias, M.M., Patz, S., and Schwartz, L.M., 1988, Magnetic resonance imaging of immiscible fluid displacement in porous media: *Physical Review Letters*, v. 61, no. 13, p. 1489-1493.
108. Chen, J.-D., and Wada, N., 1986, A new technique for visualizing the distribution of oil, water, and quartz grains in a transparent, three-dimensional, porous medium, SPE-13349: *SPE Formation Evaluation*, v. 1, no. 2, p. 205-208.
109. Chen, R., Stallworth, P.E., Greenbaum, S.G., and Kleinberg, R.L., 1994, Effects of hydrostatic pressure on proton and deuteron magnetic resonance of water in natural rock and artificial porous media: *Journal of Magnetic Resonance, Series A*, v. 110, p. 77-81.
110. Chen, Q., Wang, W., and Cai, X., 1996, Application of NMR imaging to steam foam flooding in porous media: *Magnetic Resonance Imaging*, v. 14, no. 7/8, p. 949-950.
111. Chen, S., and Georgi, D.T., 1997, Improving the accuracy of NMR relaxation distribution analysis in clay-rich reservoirs and core samples, paper SCA-9702, in 1997 international symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 10 p.
112. Chen, S., Kim, K.-H., Qin, F., and Watson, A.T., 1992, Quantitative NMR imaging of multiphase flow in porous media: *Magnetic Resonance Imaging*, v. 10, p. 815-826.
113. Chen, S., Liaw, H.K., and Watson, A.T., 1993, Fluid saturation-dependent nuclear magnetic resonance spin-lattice relaxation in porous media and pore structure analysis: *Journal of Applied Physics*, v. 74, no. 3, p. 1473-1479.
114. Chen, S., Liaw, H.-K., and Watson, A.T., 1994, Measurements and analysis of fluid saturation-dependent NMR relaxation and linebroadening in porous media: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 201-202.
115. Chen, S., Qiao, J., Liaw, H.K., and Watson, A.T., 1993, Characterization of rock pore structures using NMR restricted diffusion measurements and analyses, paper SCA-9313, in Annual technical

- conference proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 15 p.
116. Chen, S., Qin, F., Kim, K-H., and Watson, A.T., 1992, NMR imaging of multiphase flow in porous media, SPE-24760, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 1013-1026. Also published in 1993: AIChE [American Institute of Chemical Engineers] Journal, v. 39, no. 6, p. 925-934.
 117. Chen, S., Qin, F., and Watson, A.T., 1994, Determination of fluid saturations during multiphase flow experiments using NMR imaging techniques: AIChE Journal, v. 40, p. 1238-1245.
 118. Chen, S., Yao, X., Qiao, J., and Watson, A.T., 1994, Characterization of fractures and multiphase flow in fractured permeable rocks using NMR imaging techniques, paper SCA-9402, in SCA international symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts, Chapter-at-Large, p. 17-28. Also published in 1994 as, NMRI characterization of fractures and multiphase transport in fractured porous media, SPE-28369, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 65-79.
 119. Chen, S., Yao, X., Qiao, J., and Watson, A.T., 1995, Characterization of fractured permeable porous media using relaxation-weighted imaging techniques: Magnetic Resonance Imaging, v. 13, no. 4, p. 599-606.
 120. Cherry, R., 1997, Magnetic resonance technology and its applications in the oil and gas industry, part 1: Petroleum Engineer International, 70, no. 3, p. 29-35.
 121. Coates, G.R., 1993, Magnetic resonance well logging, in New technology for independent producers, conference proceedings: Rocky Mountain Association of Geologists, Denver, section 6, p. 29-41.
 122. Coates, G., Collier, H., Milligan, B., Vasilache, M., and Carter, J., 1992, MRIL, an environmentally safe measure of porosity and permeability, in 4th international symposium on borehole geophysics for minerals, geotechnical and groundwater applications, proceedings: Society of Professional Well Log Analysts, Minerals and Geotechnical Logging Society, Chapter-at-Large, p. 21-35.
 123. Coates, G.R., and Howard, A.D., 1992, Core data and the MRIL show a new approach to 'formation factor', SCA-9207, in 6th annual technical conference preprints [33rd SPWLA annual logging symposium transactions, v. 3]: Society of Core Analysts, Society of Professional Well Log Analysts Chapter-at-Large, 14 p.
 124. Coates, G.R., Gardner, J.S., and Miller, D.L., 1994, Applying pulse-echo NMR to shaly sand formation evaluation, paper B, in 35th annual logging symposium transactions: Society of Professional Well Log Analysts, 22 p.
 125. Coates, G.R., Gardner, J.S., Vinegar, H.J., and Tutunjian, P.N., 1993, Restrictive diffusion from uniform gradient NMR well logging, SPE-26472, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 575-590.
 126. Coates, G., Marschall, D., Mardon, D., and Galford, J., 1997, A new characterization of bulk-volume irreducible using magnetic resonance, paper QQ, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 14 p. Also published in 1997: DiaLog [London Petrophysical Society], v. 5, no. 6, p. 9-16.
 127. Coates, G., and Miller, D., 1997, Confirming Rw by reconstructing the SP using formation evaluation results employing gradient field magnetic resonance logs, paper VV, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p.
 128. Coates, G.R., Miller, M., Gillen, M., and Henderson, G., 1991, The MRIL in Conoco 33-1--an investigation of a new magnetic resonance imaging log, paper DD, in 32nd annual logging symposium transactions: Society of Professional Well Log Analysts, 24 p.
 129. Coates, G.R., Miller, D.L., Mardon, D., and Gardner, J.S., 1995, Applying log measurements of

- restricted diffusion and T2 to formation evaluation, paper P, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 11 p.
130. Coates, G.R., Peveraro, R.C.A., Hardwick, A., and Roberts, D., 1991, The Magnetic Resonance Imaging Log characterized by comparison with petrophysical properties and laboratory core data, SPE-22723, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 627-635.
 131. Coates, G.R., Vinegar, H.J., Tutunjian, P.N., and Gardner, J.S., 1993, Restrictive diffusion from uniform gradient NMR well logging, SPE-26472, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 575-590.
 132. Cohen, M.H., and Mendelson, K.S., 1982, Nuclear magnetic resonance and the internal geometry of sedimentary rocks: *Journal of Applied Physics*, v. 53, no. 2, p. 1127-1135.
 133. Cook, J.C., 1977, Electromagnetic resonance borehole assay logging, U.S. Bureau of Mines Open-File Report No. 69-78, 100 p.
 134. Cook, J.C., 1979, Electromagnetic resonance borehole assay logging [final report]. U.S. Bureau of Mines Open-File Report No. TR-77-10, 103 p.
 135. Coolidge, J., 1962, Nuclear magnetism logging: *Oil and Gas Journal*, v. 60, no. 10, p. 92-97.
 136. Coolidge, J.E., Ritchie, I., Gamson, B.W., and Shane, L., 1962, Recent advances in nuclear magnetism logging, paper 11, in 3rd annual logging symposium transactions: Society of Professional Well Log Analysts, 24 p.
 137. Coope, D.F., 1994, Petrophysical applications of NMR in the North Sea, paper Z, in 16th European formation evaluation symposium transactions: Society of Professional Well Log Analysts, Aberdeen Chapter, 12 p. Abridged version published also published in 1994 as, Petrophysical applications of NMR: *Dialog* [London Petrophysical Society Quarterly Newsletter], v. 2, no. 4, October, p. 8-10.
 138. Corbelleri, A., Lenge, D., Ortea, J., and Breda, E., 1996, Application of the magnetic resonance logging in San Jorge basin (Argentina), paper VV, in 37th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p.
 139. Cory, D.G., 1992, Solid state NMR imaging: *Annual Reports of NMR Spectroscopy*, v. 24, p. 87-180.
 140. Cotts, R.M., Hoch, M.J.R., Sun, T., and Markert, J.T., 1989, Pulsed field gradient stimulated echo methods for improved NMR diffusion measurements in heterogeneous systems: *Journal of Magnetic Resonance*, v. 83, p. 252-266.
 141. Cowgill, D.F., Pitman, J.K., and Seevers, D.O., 1981, NMR determination of porosity and permeability of western tight gas sands, SPE-9875, in SPE/DOE low permeability gas reservoirs symposium proceedings: Society of Petroleum Engineers, p. 437-447.
 142. Crary, S., Pellegrin, F., and Simon, B., 1997, NMR applications in the Gulf of Mexico, paper PP, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 13 p.
 143. Crosnier, Y., Citerna, J., and Gabillard, R., 1972, Specific surface area of rocks and nuclear magnetic resonance logging, paper 8 [in French], in 1st European formation evaluation symposium transactions: Society of Professional Well Log Analysts, 20 p.
 144. Crowe, B., LaTorraca, G., Dunn, K., Donovan, M., Jasek, D., and Van Dalen, S., 1997, Measuring residual oil saturation in west Texas using NMR, paper JJ, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p.
 145. Cunningham, A.B., and Jay, K.L., 1991, Field experience using the nuclear magnetic logging tool for quantifying microporosity and irreducible water saturation, paper EE, in 32nd annual logging symposium transactions: Society of Professional Well Log Analysts, 23 p.
 146. Curwen, D.W., and Molaro, C., 1995, Permeability from magnetic resonance imaging logs, paper GG, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p.
 147. D'Orazio, F., Tarczon, J.C., Halperin, W.P., Eguchi, K., and Mizusaki, T., 1989, Application of

- nuclear magnetic resonance pore structure analysis to porous silica: *Journal of Applied Physics*, v. 65, p. 742-751.
148. Dashen, R., Day, P., Kenyon, W., Straley, C., and Willemsen, J., 1987, T1-permeability correlations: *American Institute of Physics Proceedings*, v. 154, no. 1, p. 37-62.
 149. Datema, K.P., Bolt-Wsterhoff, J.A., Nesbitt, G.J., Maarsen, P.K., Ylstra, W., tutunjian, P.N., Viengar, H., and Karger, J., 1992, Petrochemical applications of pulsed field gradient NMR, in *Magnetic resonance microscopy*: VCH Publishers, New York, p. 395-416.
 150. Davies, S., Hardwick, A., Roberts, D., Spowage, K., and Packer, K.J., 1994, Quantification of oil and water in preserved reservoir rock by NMR spectroscopy and imaging: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 349-353.
 151. Davies, S., Hardwick, A., Spowage, K., and Packer, K.J., 1994, Fluid velocity imaging of reservoir and core samples: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 265-268.
 152. Davies, S., Kalam, M.Z., Packer, K.J., and Zelaya, F.O., 1990, Pore-size distributions from NMR spin-lattice relaxation measurements of fluid saturated porous solids, part 2, Application to reservoir core samples: *Journal of Applied Physics*, v. 67, no. 6, p. 3171-3176.
 153. Davies, S., and Packer, K.J., 1990, Pore-size distributions from NMR spin-lattice relaxation measurements of fluid-saturated porous solids, part 1, Theory and simulation: *Journal of Applied Physics*, v. 67, p. 3163-3170.
 154. Davies, S., Packer, K.J., Roberts, D.R., and Zelaya, F.O., 1991, Pore-size distributions from NMR spin-lattice relaxation data: *Magnetic Resonance Imaging*, v. 9, p. 681-685.
 155. Dechter, J.J., Komoroski, R.A., Ramaprasad, S. 1989, NMR chemical shift selective imaging of individual fluids in sandstone and dolomite cores, paper SCA-8903, in *Annual technical conference reprints*, v. 1 (1987-1989): Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 14 p.
 156. Dennis, B., 1997, Pores explored: *Schlumberger Middle East Well Evaluation Review*, no. 19, p. 53-65.
 157. Dereppe, J.M., and Moreaux, C., 1991, 2D spin-echo and 3D chemical-shift-imaging techniques for analysis of oil-water replacement in limestone: *Journal of Magnetic Resonance*, v. 91, p. 596-603.
 158. Dereppe, J.M., Moreaux, C., and Schenker, K., 1991, Chemical shift imaging of fluid filled porous rocks: *Magnetic Resonance Imaging*, v. 9, p. 809-813.
 159. Devereaux, O.F., 1967, Effect of crude oil on the nuclear magnetic relaxation of water protons in sandstone: *Nature*, v.215, no. 5101, p. 614-615.
 160. Dodge, W.S., Shafer, J.L., and Guzman-Garcia, A.G., 1995, Core and log NMR measurements of an iron-rich glauconitic sandstone reservoir, paper O, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p.
 161. Dodge, W.S., Shafer, J., and Klimentidis, R., 1996, Capillary pressure--the key to producible porosity, paper J, in 37th annual logging symposium transactions: Society of Professional Well Log Analysts, 13 p.
 162. Doughty, D.A., and Maerefat, N.L., 1987, Transformation of an NMR spectrometer into an NMR imager for evaluating fluid content and rock properties of core samples, paper M, SCA-8713, in *Annual technical conference proceedings*, 6 p. Later reprinted in 1989, in *Annual technical conference reprints*, v. 1, part 1: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 6 p. Later published in 1989: *The Log Analyst*, v. 30, no. 2, p. 78-84.
 163. Doughty, D.A., and Tomutsa, L., 1992, NMR microscopy for fluid imaging at pore scale in reservoir rocks, SCA-9222, in 6th annual technical conference preprints [33rd SPWLA annual logging symposium transactions, v. 3]: Society of Core Analysts, Society of Professional Well Log Analysts Chapter-at-Large, 15 p.
 164. Doughty, D.A., and Tomutsa, L., 1993, Investigation of wettability by NMR microscopy and spin-lattice relaxation: U.S. Department of Energy Report NIPER-720, 11 p.
 165. Doughty, D.A., and Tomutsa, L., 1996, Multinuclear NMR spectroscopy of two-phase fluid

- systems in porous rock: *Magnetic Resonance Imaging*, v. 14, no. 7/8, p. 869-873.
166. Duda, L.E., and Pitman, J.K., 1982, Pore structure measurements, in *Western tight gas sands advanced logging workshop: U.S. Department of Energy Report CONF-8103113*, p. 98-127.
 167. Dunn, K-J., LaTorraca, G.A., Warner, J.L., and Bergman, D.J., 1994, On the calculation and interpretation of NMR relaxation time distributions, SPE-28367, in *SPE annual technical conference and exhibition proceedings*, v. omega, *Formation evaluation and reservoir geology: Society of Petroleum Engineers*, p. 45-54.
 168. Edelstein, W.A., 1991, Oil core NMR imaging/spectroscopy instrumentation: *Magnetic Resonance Imaging*, v. 9, no. 5, p. 865-868.
 169. Edelstein, W.A., Roemer, P.B., Mueller, O.M., Vinegar, H.J., and Tutunjian, P.N., 1988, NMR imaging for core analysis, SPE-18272, in *1988 SPE annual technical conference and exhibition proceedings*, v. sigma, *Reservoir engineering: Society of Petroleum Engineers*, p. 101-112.
 170. Edelstein, W.A., Vinegar, H.J., Tutunjian, P.N., and Roemer, P.B., 1990, Time-domain curve fitting for automated NMR spectral analysis, paper 9025, in *Annual technical conference preprints*, v. 2: *Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large*, 17 p.
 171. Edwards, C., 1997, Effects of tool design and logging speed on T2 NMR log data, paper RR, in *38th annual logging symposium transactions: Society of Professional Well Log Analysts*, 13 p.
 172. Edwards, C., and Chen, S., 1996, Improved NMR well logs from time-dependent echo filtering, paper RR, in *37th annual logging symposium transactions: Society of Professional Well Log Analysts*, 12 p.
 173. Egbogah, E.O., Amar, Z.H.B.T. Amar, 1997, Accurate initial/residual saturation determination reduces uncertainty in further development and reservoir management of the Duland field, offshore peninsular Malaysia, SPE-38024, in *Asia Pacific oil and gas conference proceedings: Society of Petroleum Engineers*, p. 79-90.
 174. Eidesmo, T., Relling, O., and Rueslatten, H., 1996, NMR logging of mixed-wet North Sea sandstone reservoirs, paper PP, in *37th annual logging symposium transactions: Society of Professional Well Log Analysts*, 14 p.
 175. Ellis, D.V., 1987, Nuclear magnetic logging, chapter 14, in *Well logging for earth scientists: Elsevier Science Publishing Company, New York*, p. 305-337.
 176. Enwere, M.P., and Archer, J.S., 1992, NMR imaging for core flood testing, SCA-9218, in *6th annual technical conference preprints [33rd SPWLA annual logging symposium transactions, v. 3]: Society of Core Analysts, Society of Professional Well Log Analysts Chapter-at-Large*, 14 p.
 177. Enwere, M.P., and Archer, J.S., 1992, NMR imaging for water/oil displacement in cores under viscous-capillary force control, SPE-24166, in *8th Enhanced oil recovery symposium proceedings: Society of Petroleum Engineers*, p. 99-104.
 178. Fatt, I., Kumar, J., and Saraf, D.N., 1969, Nuclear magnetic relaxation time of water in a porous medium with heterogeneous surface wettability: *Journal of Applied Physics*, v. 40, no. 10, p. 4165-4171.
 179. Fertl, W.H., 1984, Recent advances in the evaluation of heavy oil reservoirs based on geophysical well-logging concepts, in R.F. Meyer, J.C. Wynn and J.C. Olson, eds., *The future of heavy crude and tar sands [2nd international UNITAR conference proceedings, 1982]: McGraw-Hill Inc., Coal Age Mineral Information Service, New York*, p. 313-317.
 180. Fertl, W.H., 1987, Well log assisted evaluation of heavy-oil resources; a status report, in R.F. Myer, ed., *Exploration for heavy crude oil and natural bitumen [AAPG/UNITAR research conference (Santa Maria, California, October 29-November 2, 1984), proceedings]: AAPG Studies in Geology No. 25*, p. 443-452.
 181. Flaum, C., Kleinberg, R., and Hurlimann, M., 1996, Identification of gas with the Combinable Magnetic Resonance Tool (CMR), paper L, in *37th annual logging symposium transactions: Society of Professional Well Log Analysts*, 11 p.
 182. Foley, Farooqui, S.A., and Kleinberg, R.L., 1996, NMR relaxation of fluids at solid surfaces--

- effects of paramagnetic ions: *Journal of Magnetic Resonance, Series A*, v. 123, p. 95.
183. Fordham, E.J., Gibbs, S.J., and Hall, L.D., 1994, Partially restricted diffusion in a permeable sandstone--observations by stimulated echo PFG NMR: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 279-284.
 184. Fordham, E.J., Hall, L.D., and Sezginer, A., 1995, Imaging multiexponential relaxation in the (y , $\log(e) T_1$) plane, with application to clay filtration in rock cores: *Journal of Magnetic Resonance, Series A*, v. 113, p. 139-150.
 185. Fordham, E.J., Horsfield, M.A., Hall, C., and Hall, L.D., 1991, Low-contrast secondary imbibition in long rock cores: *Magnetic Resonance Imaging*, v. 9, p. 803-808.
 186. Fordham, E.J., Horsfield, M.A., Hall, L.D., and Maitland, G.C., 1993, Depth filtration of clay in rock cores observed by one-dimensional 1H NMR imaging: *Journal of Colloid and Interface Science*, v. 156, p. 253-255.
 187. Fordham, E.J., Roberts, T.P.L., Carpenter, T.A., Hall, L.D., and Hall, C., 1991, Dynamic NMR imaging of rapid depth filtration of clay in porous media: *AIChE [American Institute of Chemical Engineering] Journal*, v. 37, no. 12, p. 1900-1903.
 188. Foster, M.A., and Hutchinson, J.M.S., 1987, *Practical NMR imaging*: ILR, Oxford Press, Washington, D.C.
 189. Freedman, R., and Morriss, C.E., 1995, Processing data from an NMR logging tool, SPE-30560, in *Annual technical conference and exhibition proceedings*, v. omega, *Formation evaluation and reservoir geology*: Society of Petroleum Engineers, p. 301-316.
 190. Freedman, R., and Rouault, G.F., 1987, Nuclear magnetism logging; signal processing for residual oil saturation, SPE-16771, in *SPE annual technical conference and exhibition, proceedings*, v. omega, *Formation evaluation and reservoir geology*: Society of Petroleum Engineers, p. 309-321. Later published in 1989 as, *Remaining-oil determination using nuclear magnetism logging*: *SPE Formation Evaluation*, v. 4, no. 2, p. 21-130.
 191. Freedman, R., Boyd, A., Gubelin, G., McKeon, D., Morriss, C., and Flaum, C., 1997, Measurement of total NMR porosity adds new value to NMR logging, paper OO, in *38th annual logging symposium transactions*: Society of Professional Well Log Analysts, 14 p.
 192. Fukushima, E., and Roeder, S.B.W., 1981, *Experimental pulse NMR--a nuts and bolts approach*: Addison-Wesley, Reading, Massachusetts.
 193. Gaigala, A.K., Chidester van Orden, A., Robertson, B., Mareci, T., and Lewis, L.A., 1989, Application of magnetic resonance imaging to visualization of flow in porous media: *Nuclear Technology*, v. 84, p. 113-118.
 194. Gallegos, D.P., Munn, K., Douglas, M.S., and Stermer, D.L., 1988, NMR technique for the analysis of pore structure application to materials with well-defined pore structure: *Journal of Colloid and Interface Science*, v. 119, no. 1, p. 127-140.
 195. Gallegos, D.P., and Smith, D.M., 1988, A NMR technique for the analysis of pore structure--determination of continuous pore size distributions: *Journal of Colloid and Interface Science*, v. 122, no. 1, p. 143-153.
 196. Gallegos, D.P., Smith, D.M., and Brinker, C.J., 1988, An NMR technique for the analysis of pore structure--application to mesopores and micropores: *Journal of Colloid and Interface Science*, v. 124, p. 186-198.
 197. Gamson, B.W., Stephenson, T.B., Edwards, L., and Shane, L., 1963, Recent nuclear magnetism logging field results in Texas and Oklahoma: *The Log Analyst*, v. 2, no. 2, p. 1-58.
 198. Geissler, E., and Hecht, A.M., 1970, Nuclear magnetic resonance and relaxation of adsorbed water in synthetic fluor montmorillonite: *Journal of Colloid and Interface Science*, v. 34, no. 1, p. 32-35.
 199. Georgi, D.T., editor, 1995, *Nuclear magnetic resonance logging short course notes*, 36th annual symposium, Paris, France: Society of Professional Well Log Analysts, variously paginated.
 200. Georgi, D.T., Cullen, M.P., Buchanan, T.R., Huber, P.M., and Lyon, S.A., 1991, *Formation evaluation contributions to the Cessford shallow gas pilot*, paper F, in *13th Formation evaluation*

- symposium transactions: Canadian Well Logging Society, 27 p.
201. Georgi, D.T., Zhang, J.C. Huang, Y.Y., Chandler, R.N. and Prammer, M.G., 1995, Vertical resolution and signal-to-noise characteristics of nuclear magnetic resonance imaging logs, paper OOO, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p.
 202. Gerritsma, C.J., Oosting, P.H., and Trappeniers, N.J., 1971, Proton spin-lattice relaxation and self-diffusion in methane, part 2--Experimental results for proton spin-lattice relaxation times: *Physica*, v. 51, p. 381-394.
 203. Gerritsma, C.J., and Trappeniers, N.J., 1971, Proton spin-lattice relaxation and self-diffusion in methanes, part 1, Spin-echo spectrometer and preparation of the methane samples: *Physica*, v. 51, p. 365-380.
 204. Gibbs, S.J., Attard, J.J., and Hall, L.D., 1993, Spatially determined water diffusivities in brine saturated reservoir cores obtained by NMR imaging: *AIChE Journal*, v. 39, p. 689-694.
 205. Gibson, B-D., Boyd, A., Poon, A., and Gubelin, W.G., 1995, Nuclear magnetic resonance applications in North America, paper H, in 1st annual well logging symposium of Japan, proceedings: Society of Professional Well Log Analysts, Japan Chapter, 14 p.
 206. Gladden, L.F., 1993, Nuclear magnetic resonance studies of porous media: *Chemical Engineering Research Descriptions* [?], v. 71, p. 657-674.
 207. Gladden, L.F., 1994, Nuclear magnetic resonance in chemical engineering--principles and applications: *Chemical Engineering Science*, v. 49, no. 20, p. 3339-3408.
 208. Glasel, J.A., 1970, NMR relaxation in heterogeneous systems: *Nature*, v. 227, p. 704-705.
 209. Glasel, J.A., and Lee, K.H., 1979, On the interpretation of water nuclear magnetic resonance relaxation times in heterogeneous systems: *Journal of the American Chemical Society*, v. 96, p. 970-978.
 210. Gleeson, J.W., and Woessner, D.E., 1991, Three-dimensional and flow-weighted NMR imaging of pore connectivity in a limestone: *Magnetic Resonance Imaging*, v. 9, p. 879-884.
 211. Gleeson, J.W., Woessner, D.W., and Jordan, C.F., Jr., 1990, NMR imaging of pore structures in limestones, SPE-20493, in Annual technical conference proceedings, v. gamma, EOR/general petroleum engineering: Society of Petroleum Engineers, p. 247-253. Later published in 1993: *SPE Formation Evaluation*, v. 6, no. 2, p. 123-127.
 212. Goelman, G., and Prammer, M.G., 1995, The CMPG pulse sequence in strong magnetic field gradients with applications to oil-well logging: *Journal of Magnetic Resonance, Series A*, v. 113, p. 11-18.
 213. Gogolashvili, E., 1996, Wettability and T1 proton relaxation times of sandstone rocks: *Magnetic Resonance Imaging*, v. 14, no. 7/8, p. 1003-1005.
 214. Gonfalini, M., and Chelini, V., 1994, High technology logging of complex reservoirs in the Adria project, in 14th World petroleum congress proceedings, v. 2, Exploration, production, and reserves: John Wiley and Sons, New York, p. 133-134.
 215. Goodman, B.A., and Chudek, J.A., 1994, Nuclear magnetic resonance spectroscopy, chapter 4, in Wilson, M.J., editor, *Clay mineralogy--spectroscopic and chemical determinative methods*: Chapman and Hall, London, p 120-172.
 216. Gossenberg, P., Galli, G., Andreani, M., and Klopff, W., 1996, A new petrophysical interpretation model for clastic rocks based on NMR, epithermal neutron and electromagnetic logs, paper M, in 37th annual logging symposium transactions: Society of Professional Well Log Analysts, 13 p.
 217. Griffin, D.D., Kleinberg, R.L., and Fukuhara, M., 1993, Low frequency NMR spectrometer: *Measurement Science and Technology*, v. 4, p. 968.
 218. Gronas, T., Roaldset, E., Rueslatten, H., Skjetne, T., and Roaldset, E., 1995, NMR used to study pore sizes in sand, F028, in 57th EAEG Conference and Technical Exhibition Extended Abstracts, v. 2: European Association of Exploration Geophysicists, 2 p.
 219. Gronas, T., Rueslatten, H., Roaldset, E., and Skjetne, T., 1996, NMR responses to kaolinite in sand: *Magnetic Resonance Imaging*, v. 14, no. 7/8, p. 961-962.

220. Gubelin, G., and Boyd, A., Total porosity and bound-fluid measurements from an NMR tool, SPE-39096: Journal of Petroleum Technology, v. 49, no. 7, p. 718.
221. Guilfoyle, D.N., and Mansfield, P., 1991, MRI in porous media--flow measurement in porous media by echo-planar imaging: Magnetic Resonance Imaging, v. 9, no. 5, p. 775-777.
222. Guilfoyle, D.N., Mansfield, P., and Packer, K.J., 1992, Fluid flow measurement in porous media by echo-planar imaging: Journal of Magnetic Resonance, v. 97, p. 342-358.
223. Guillen, A., 1997, Inverse problem of magnetic resonance measurements applied to water resource characterization, EM 4.3, in 1997 technical program expanded abstracts with authors' biographies, v. 1: Society of Exploration Geophysicists, Tulsa, p. 446-449.
224. Guillot, G., 1988, Magnetic resonance imaging of fluids in porous media--a comparison to other experimental techniques, in Fuillon, E., and others, editors, Disorder and mixing: Kluwer, Dordrecht, p. 319-326.
225. Guillot, G., Chardaire-Riviere, C., Bobroff, S., Le Roux, A., Roussel, J.C., and Cuiec, L., 1994, Characterisation of wetting heterogeneities in sandstone rocks by MRI: Magnetic Resonance Imaging, v. 12, no. 2, p. 365-368.
226. Guillot, G., Trokiner, A., Darasse, L., and Saint-Jalmes, H., 1989, Drying of a porous rock monitored by NMR imaging: Journal of Physics, Series D, Applied Physics, v. 22, p. 1646-1649.
227. Guillot, G., Trokiner, A., Darasse, L., Dupas, A., Ferdossi, F., Kassab, G., Hulin, J.P., Rigord, P., and Saint-Jalmes, H., 1991, NMR imaging applied to various studies of porous media: Magnetic Resonance Imaging, v. 9, p. 821-825.
228. Gumilar, B., Adriansyah, R., Thomas, A.R., and Darmawan, B., 1996, Signal processing method for multiexponentially decaying signals and an analysis of low-contrast pay in Telisa sands packages in central Sumatra, paper IPA96-2-2-123, in 25th annual convention proceedings, v. 2: Indonesia Petroleum Association, Jakarta, p. 175-187.
229. Gummerson, R.J., Hall, C., and Hoff, W.D., 1979, Unsaturated water flow within porous materials observed by NMR imaging: Nature, v. 281, p. 56-57.
230. Hall, L.D., and Rajanayagam, V., 1987, Thin-slice, chemical-shift imaging of oil and water in sandstone rock at 80 Mhz: Journal of Magnetic Resonance, v. 74, p. 139-146.
231. Hall, L.D., Rajanayagam, V., and Hall, C., 1986, Chemical-shift imaging of water and n-dodecane in sedimentary rocks: Journal of Magnetic Resonance, v. 68, p. 185-188.
232. Halperin, W.P., Bhattacharya, S., and D'Orazio, F.D., 1991, Relaxation and dynamical properties of water in partially filled porous materials using NMR techniques: Magnetic Resonance Imaging, v. 9, po. 733-737.
233. Halperin, W.P., D'Orazio, F.D., Bhattacharya, S., and Tarczon, J.C., 1989, Magnetic resonance relaxation in porous media, chapter 11, in Klafter, J., and Drake, J.M., eds., Molecular dynamics in restricted geometries: Wiley Interscience Publications, New York, p. 311-350.
234. Harben, P.E., 1987, Nuclear magnetic resonance imaging and spectroscopy: Lawrence Livermore National Laboratory Report No. UCRL-50025-87-2, 8 p.
235. Hartman, D.E., Loren, J.D., Robinson, J.D., and Vajnar, E.A., 1972, Determination of residual oil using the nuclear magnetism log, SPE-3797, in Improved oil recovery symposium preprints: Society of Petroleum Engineers, Mid-Continent Section, p. 263-280.
236. Hassoun, T.H., Zainalebedin, K., and Minh, C.C., 1997, Hydrocarbon detection in low-contrast resistivity pay zones, capillary pressure and ROS determination with NMR logging in Saudi Arabia, SPE-37770, in 10th SPE Middle East oil show conference proceedings, v. 2: Society of Petroleum Engineers, p. 127-143.
237. Hatcher, P.G., and Romankiw, L.A., 1985, Nuclear magnetic resonance studies of organic-matter-rich sedimentary rocks of some early Mesozoic basins of the eastern United States: U.S. Geological Survey Circular No. 946, p. 65-70.
238. Hayward, R.J., Packer, K.J., and Tomlinson, D.J., 1972, Pulsed field-gradient spin-echo NMR studies of flow in fluids: Molecular Physics, v. 23, p. 1083.
239. Hazlett, R.D., Furr, M.J., and Navarro, R., 1992, A correlation for miscible flood displacement

- efficiency in the San Andres with NMR relaxation, SPE-23992, in Permian Basin oil and gas recovery conference proceedings: Society of Petroleum Engineers. Later published in 1994: SPE Advanced Technology Series, v.2, no. 2, p. 185-190.
240. Hazlett, R.D., Gleeson, J.W., Laali, H., and Navarro, R., 1993, NMR imaging application to CO₂ miscible flooding in west Texas carbonates, paper SCA-9311, in Annual technical conference proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 9 p.
241. Headley, L.C., 1973, NMR relaxation of ⁷Li and ¹H in Appalachian petroleum reservoir rocks containing LiCl solution: *Nature*, v. 242, no. 119, April 9, p. 87-88.
242. Hedberg, S.A., Knight, R.J., MacKay, A.L., and Whittall, K.P., 1993, The use of nuclear magnetic resonance for studying and detecting hydrocarbon contaminants in porous rock: *Water Resources Research*, v. 29, no. 4, p. 1163-1170.
243. Hemminga, M.A., and de Jager, P.A., and Sonneveld, A., 1977, The study of flow by pulsed nuclear magnetic resonance, part 2, Measurement of flow velocities using a repetitive pulse method: *Journal of Magnetic Resonance*, v. 37, p. 1-16.
244. Herrick, R.C., Couturie, S.H., and Best, D.L., 1979, An improved nuclear magnetism logging system and its application to formation evaluation, SPE-8361: Society of Petroleum Engineers, 54th Annual Technical Conference and Exhibition Preprints, 7 p.
245. Hilchie, D.W., 1988, NML--The hope for production prediction: *Journal of Petroleum Technology*, v. 40, no. 3, P. 273-275.
246. Hinedi, Z.R., Chang, A.C., Anderseon, M.A. and Borchardt, D.B., 1997, Quantification of microporosity by nuclear magnetic resonance relaxation of water imbibed in porous media: *Water Resources Research*, v. 33, no. 12, p. 2697.
247. Holt, R.M., Fjaer, E., Torsaeter, O., and Bakke, S., 1996, Petrophysical laboratory measurements for basin and reservoir evaluation: *Marine and Petroleum Geology*, v. 13, no. 4, p. 383-391.
248. Hook, J., Basan, P., and Lowden, B., 1996, The MRIL log in a low porosity and permeability formation, paper N, in 37th annual logging symposium transactions: Society of Professional Well Log Analysts, 13 p.
249. Horkowitz, J.P., Hartman, D.E., Clerke, E.A., Vinegar, H.J., and Coates, G.R., 1995, Residual oil saturation measurements in carbonates with pulsed NMR logs, paper Q, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p. Later published in 1997: *The Log Analyst*, v. 38, no. 2, p. 73-83.
250. Hom, G.M., 1996, The Central Graben--a dynamic overpressure system, in Glennie, K., and Hurst, A., eds., AD1995--NW Europe's hydrocarbon industry: Geological Society [London], p. 177-184.
251. Horsfield, M., Hall, C., and Hall, L.D., 1990, Two-species chemical shift imaging using prior knowledge and estimation theory, application to rock cores: *Journal of Magnetic Resonance*, v. 87, p. 319-330.
252. House, W.V., 1983, Theoretical basis for NMR imaging, in Partin, C.L., James, A.E., Rollo, G.D., and Price, R.R., eds., *Nuclear magnetic resonance NMR imaging*: W.B. Saunders Company, Philadelphia, Pennsylvania, p. 60-72.
253. Howard, A., 1996, MRI logging tools locate water-free production: *American Oil and Gas Reporter*, v. 39, no. 11, October, p. 70, 72-74.
254. Howard, J.J., 1994, Wettability and fluid saturations determined from NMR T₁ distributions: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 197-200.
255. Howard, J.J., and Kenyon, W.E., 1992, Determination of pore size distribution in sedimentary rocks by proton nuclear magnetic resonance: *Marine and Petroleum Geology*, v. 9, no. 2, p. 139-145.
256. Howard, J.J., Kenyon, W.E., and Straley, C., 1990, Proton-magnetic-resonance and pore-size variations in reservoir sandstones, SPE-20600, in SPE annual technical conference exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum

- Engineers, p. 733-742. Later published in 1993: SPE Formation Evaluation, v. 8, no. 3, p. 194-200.
257. Howard, J.J., and Spinler, E.A., 1993, Nuclear magnetic resonance measurements of wettability and fluid saturations in chalk, SPE-26471, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 565-573. Later published in 1995: SPE Advanced Technology Series, v. 3, no. 1, p. 60-65.
258. Howard, J., Williams, J., and Thorpe, D., 1997, Permeability from nuclear magnetic resonance logging in a gas-condensate field, paper XX, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 13 p.
259. Hsu, W.F., 1994, Wettability of porous media by nuclear magnetic resonance relaxation methods: Texas A&M University, College Station, Texas, unpublished Ph.D. thesis, 159 p.
260. Hsu, W-F., Li, X., and Flumerfelt, R.W., 1992, Wettability of porous media by NMR relaxation methods, SPE-24761, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 1027-1037.
261. Hull, P.W., and Coolidge, J.E., 1960, Field results with nuclear magnetism logging, paper 11, in 1st annual logging symposium transactions: Society of Professional Well Log Analysts, 26 p.
262. Hurlimann, M.D., Helmer, K.G., Latour, L.L., and Sotak, C.H., 1994, Restricted diffusion in sedimentary rocks--determination of surface-area-to-volume ratio and surface relaxivity: Journal of Magnetic Resonance, Series A, v. 111, p. 169.
263. Hurlimann, M.D., Latour, L.L., and Sotak, C.H., 1994, Diffusion measurement in sandstone core--NMR determination of surface-to-volume ratio and surface relaxivity: Magnetic Resonance Imaging, v. 12, p. 325-327.
264. Issa, B., and Mansfield, P., 1994, Permeability estimation from T1 mapping: Magnetic Resonance Imaging, v. 12, no. 2, p. 213-214.
265. Jackson, J.A., 1981, New NMR well logging/fracture mapping technique with possible application of Squid NMR detection, in Weinstock, H., and Overton, W.C., Jr., editors, Workshop on Squid applications to geophysics: Society of Exploration Geophysicists, p. 161-165.
266. Jackson, J.A., 1984, Nuclear magnetic resonance logging: The Log Analyst, v. 25, no. 5, September-October, p. 16.
267. Jackson, J.A., Brown, J.A., and Crawford, T.R., 1981, Remote characterization of tight gas formations with a new NMR logging tool: Los Alamos National Laboratory Report No. LA-UR-81-735, 18 p. Also published in 1981 as SPE/DOE-9860, in 1981 SPE/DOE symposium on low permeability gas reservoirs, proceedings: Society of Petroleum Engineers, p. 313-320.
268. Jackson, J.A., Brown, J.A., and Koelle, A.R., 1982, Western gas sands project, Los Alamos NMR (nuclear magnetic resonance) well logging tool development, progress report, April 1, 1980-September 30, 1981: Los Alamos National Laboratory Report No. LA-9151-PR, 24 p. Also published in 1982: U.S. Department of Energy Fossil Energy Report No. DOE/BC/10018-20, 34 p.
269. Jackson, J.A., Brown, J.A., and Koelle, A.R., 1982, Western gas sands project, Los Alamos NMR (nuclear magnetic resonance) well logging tool development, progress report, October 1, 1981-September 30, 1982: Los Alamos National Laboratory Report No LA-9641-PR, 30 p.
270. Jackson, J., and Mathews, M., 1993, Nuclear magnetic resonance bibliography: The Log Analyst, v. 34, no. 3, p. 35-69.
271. Jackson, S.R., 1993, Reservoir assessment and characterization: U.S. Department of Energy Report NIPER-685, p. 17-101.
272. Jerosch-Herold, M., Thomann, H., and Thompson, A.H., 1991, Nuclear magnetic resonance relaxation in porous media, SPE-22861, in Annual technical conference and exhibition proceedings: Society of Petroleum Engineers.
273. Jerosch-Herold, M., Thomann, H., and Thompson, A.H., 1994, Combined proton T1N and CPMG T2N studies of water saturated sandstone core plugs: Magnetic Resonance Imaging, v. 12, no. 2,

- p. 369-373.
274. Karger, J., and Pfeifer, H., 1987, NMR self-diffusion studies in zeolite science and technology: *Zeolites*, v. 7, p. 90-107.
 275. Karger, J., Pfeifer, H., and Heink, W., 1988, Principles and application of self-diffusion measurements by nuclear magnetic resonance, in Waugh, J.S., editor, *Advances in magnetic resonance*, v. 12: Academic Press, New York, p. 1-89.
 276. Karger, J., and Ruthven, D.M., 1992, *Diffusion in zeolites and other microporous solids*: Wiley, New York.
 277. Katz, A.J., and Thompson, A.H., 1986, Quantitative prediction of permeability in porous rock: *Physical Review, Series B*, v. 34, p. 8179-8191.
 278. Kenyon, W.E., 1992, Nuclear magnetic resonance as a petrophysical measurement : *Nuclear Geophysics*, v. 6, no. 2, p. 153-171. Later revised and published in 1997 as *Petrophysical principles of applications of NMR logging: The Log Analyst*, v. 38, no. 2, p. 21-43.
 279. Kenyon, W.E., Day, P.I., Straley, C., and Willemsen, J.F., 1986, Compact and consistent representation of rock NMR data from permeability estimation, SPE-15643: Society of Petroleum Engineers, 61st annual meeting preprint, 22 p. Later published in 1988 as, A three-part study of NMR longitudinal relaxation properties of water-saturated sandstones: *SPE Formation Evaluation*, v. 3, no. 3, p. 622-636.
 280. Kenyon, W.E., Howard, J.J., Sezginer, A., Straley, C., Matteson, A., Horkowitz, K., and Ehrlich, R., 1989, Pore-size distribution and NMR in microporous cherty sandstones, paper LL, in, 30th annual logging symposium transactions: Society of Professional Well Log Analysts, 24 p.
 281. Kenyon, W., Kleinberg, R., Straley, C., Gubelin, G., and Morriss, C., 1995, Nuclear magnetic resonance imaging--technology for the 21st century: *Schlumberger Oilfield Review*, v. 7, no. 3, p. 19-33.
 282. Kenyon, W.E., and Kolleeny, J.A., 1995, NMR surface relaxivity of calcite with adsorbed Mn²⁺: *Journal of Colloid and Interface Science*, v. 170, p. 502-514.
 283. Kenyon, W.E., Takezaki, H., Straley, C., Sen, P.N., Herron, M., Matteson, A., Petricola, M.J., 1995, A laboratory study of nuclear magnetic resonance relaxation and its relation to depositional texture and petrophysical properties--carbonate Thamama Group, Mubarratz field, Abu Dhabi, SPE-29886, in 9th Middle East oil show and conference proceedings: Society of Petroleum Engineers, v. 2, p. 477-502.:
 284. Kim, K.-H., Chen, S., Qin, F., and Watson, A.T., 1992, Use of NMR imaging for determining fluid saturation distributions during multiphase displacement in porous media, SCA-9219, in 6th annual technical conference preprints [33rd SPWLA annual logging symposium transactions, v. 3]: Society of Core Analysts, Society of Professional Well Log Analysts Chapter-at-Large, 18 p.
 285. Klein, J., Martin, P., and Widjanarko, W., 1997, Log water saturation model validation using NMR log and core data, paper TT, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 13 p.
 286. Kleinberg, R.L., 1994, Pore size distributions, pore coupling, and transverse relaxation spectra of porous rocks: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 271-274.
 287. Kleinberg, R.L., 1995, Petrophysics of the nuclear magnetic resonance tool, chapter 5, in Georgi, D.T., ed., *Nuclear magnetic resonance logging short course notes*, 36th annual symposium, Paris, France: Society of Professional Well Log Analysts, variously paginated.
 288. Kleinberg, R.L., 1996, Probing oil wells with NMR: *Industrial Physicist*, v. 2, no. 2, p. 18.
 289. Kleinberg, R.L., 1996, Utility of NMR T2 distributions, connection with capillary pressure, clay effect, and determination of the surface relaxivity parameter r₂: *Magnetic Resonance Imaging*, v. 14, p. 761-767.
 290. Kleinberg, R.L., 1996, Well logging, in *Encyclopedia of nuclear magnetic resonance*: John Wiley and Sons, v. 6, p. 4960-4969.
 291. Kleinberg, R.L., and Boyd, A., 1997, Tapered cutoffs for magnetic resonance bound water volume, SPE-38737, in Annual technical conference and exhibition proceedings, v. omega,

- Formation evaluation and reservoir geology, part 2: Society of Petroleum Engineers, p. 197-202.
292. Kleinberg, R.L., Farooqui, S.A., and Horsfield, M.A., 1993, T1/T2 ratio and frequency dependence of NMR relaxation in porous sedimentary rocks: *Journal of Colloid and Interface Science*, v. 158, no. 1, p. 195-198.
 293. Kleinberg, R.L., and Horsfield, M.A., 1990, Transverse relaxation processes in porous sedimentary rock: *Journal of Magnetic Resonance*, v. 88, no. 1, p. 9-19.
 294. Kleinberg, R.L., Kenyon, W.E., and Mitra, P.P., 1994, Mechanism of NMR relaxation of fluids in rock: *Journal of Magnetic Resonance, Series A*, v. 108, no. 2, p. 206-214.
 295. Kleinberg, R.L., Sezginer, A., Griffin, D.D., and Fukuhara, M., 1992, Novel NMR apparatus for investigating an external sample: *Journal of Magnetic Resonance*, v. 97, p. 466-485.
 296. Kleinberg, R.L., Straley, C., Kenyon, W.E., Akkurt, R., and Farooqui, S.A., 1993, Nuclear magnetic resonance of rocks--T1 vs. T2, SPE-26470, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 553-563.
 297. Kleinberg, R.L., and Vinegar, H.J., 1996, NMR properties of reservoir fluids: *The Log Analyst*, v. 37, no. 6, p. 20-32.
 298. Komorowski, P., McDonald, J., and Scribbins, B., 1989, Evaluation of a formation flooded by fresh water using geochemical, dielectric and nuclear magnetic measurements, paper M, in 12th Formation evaluation symposium transactions: Canadian Well Logging Society, 29 p.
 299. Kubica, P., 1995, Statistical tests of permeability estimates based on NMR measurements, paper VVV, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 11 p.
 300. Kulkarni, R.N., Watson, A.T., Nordtvedt, J.E., Brancolini, A., and Johnsen, O., 1996, Estimation of multiphase flow functions from dynamic displacement data--applications of NMR imaging, SPE-36855, in SPE European petroleum conference proceedings, v. 1: Society of Petroleum Engineers, p. 357-362.
 301. Kumar, A.T.A., Majors, P., and Rossen, W., 1995, Measurement of aperture and multiphase flow in fractures with NMR imaging, SPE-30558, in SPE annual technical conference and exhibition proceedings: Society of Petroleum Engineers, p. 279-290.. Later published in 1997: SPE Formation Evaluation, v. 12, no. 2, p. 101-107.
 302. Kumar, J., and Fatt, I., 1970, Nuclear magnetic resonance study of porosity, permeability and surface area of unconsolidated porous materials: *Log Analyst*, v. 11, no. 1, p. 13-16.
 303. Kumar, J., Fatt, I., and Saraf, D.N., 1969, Nuclear magnetic relaxation time of water in a porous medium with heterogenous surface wettability: *Journal of Applied Physics*, v. 40, p. 4165-4171.
 304. Lahajnar, G., Sobol, W.T., Lasic, D.D., Reeves, L.W., and Pintar, M.M., 1988, Bridge water diffusion in oil sands by NMR pulsed gradient proton spin-echo: *Journal of Canadian Petroleum Technology*, v. 27, no. 6, p. 133.
 305. Lamrous, O., Houi, D., Zarcone, C., and Pradere, J., 1989, Magnetic resonance imaging application to study of porous media: *Review of Applied Physics*, v. 24, no. 5, p. 607-612.
 306. Latif, A.A., Sungur, D., Stockden, I., and Minh, C.C., 1997, Borehole nuclear magnetic resonance--experience and reservoir applications in west Kuwait carbonate reservoirs, SPE-37771, in Middle East oil show proceedings, v. 2: Society of Petroleum Engineers, p. 145-154.
 307. LaTorraca, G.A. Dunn, K.J., and Bergman, D.J., 1995, Magnetic susceptibility contrast effects on NMR T2 logging, paper JJ, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 8 p.
 308. LaTorraca, G.A., Dunn, K.J., and Brown, R.J.S., 1993, Predicting permeability from nuclear magnetic resonance and electrical properties measurements, paper SCA-9312, in Annual technical conference proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 12 p.
 309. Latour, L.L., Kleinberg, R.L., and Mitra, P.P., and Sotak, C.H., 1995, Pore size distributions and tortuosity in heterogeneous porous media: *Journal of Magnetic Resonance, Series A*, 112, p. 83.

310. Latour, L.L., Kleinberg, R.L., and Sezginer, A., 1992, Nuclear magnetic resonance properties of rocks at elevated temperatures: *Journal of Colloid and Interface Science*, v. 150, p. 535.
311. Latour, L.L., Li, L., and Sotak, C.H., 1993, Improved PFG stimulated echo methods for the measurement of diffusion in inhomogeneous fields: *Journal of Magnetic Resonance, Series B*, v. 101, p. 72-77.
312. Latour, L.L., Mitra, P.P., Kleinberg, R.L., and Sotak, C.H., 1993, Time dependent diffusion coefficient of fluids in porous media as a probe of surface to volume ratio: *Journal of Magnetic Resonance, Series A*, v. 101, p. 342-346.
313. Letellier, M., Tinet, D., Maggion, R., and Fripiat, J., 1991, Study of relaxation mechanisms in clay/water systems--Determination of the surface area; application to cements: *Magnetic Resonance Imaging*, v. 9, p. 709-716.
314. Lewis, J., Williams, A., Enwere, P.M., Archer, J.S., and Taylor, D.G., 1990, Applications of magnetic resonance imaging in special core analysis studies, in Worthington, P.F., ed., *Advances in core evaluation, accuracy, and precision in reserves estimation [Reviewed proceedings of the first Society of Core Analysts European core analysis symposium, May 21-23, London, UK]*: Gordon and Breach Science Publishers, New York, p. 317-345.
315. Liaw, H.-K., Kulkarni, R., Chen, S., and Watson, A.T., 1994, The use of magnetic resonance spectroscopy to probe microscopic structures and fluid distributions in porous media, paper SCA-9403, in *SCA international symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts, Chapter-at-Large*, p. 29-42.
316. Liaw, H.-K., Kulkarni, R., Chen, S., and Watson, A.T., 1996, Characterization of fluid distributions in porous media by NMR techniques: *AIChE [American Institute of Chemical Engineers] Journal*, v. 42, no. 2, p. 538-546.
317. Lipsicas, M., Banavar, J.R., and Willemsen, J., 1986, Surface relaxation and pore sizes in rocks--a nuclear magnetic resonance analysis: *Applied Physics Letters*, v. 48, no. 22, p. 1544-1546.
318. Logan, W.D., Horkowitz, J.P., Laronga, R., and Cromwell, D., 1997, Practical application of NMR logging in carbonate reservoirs, SPE-38740, in *Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology, part 2: Society of Petroleum Engineers*, p. 203-215.
319. Logan, D., Morriss, C., and Williams, R., 1996, Comparison of nuclear magnetic resonance techniques in the Delaware, SPE-30627, in *Permian Basin oil and gas recovery conference proceedings: Society of Petroleum Engineers*, p. 75-90. Also published in 1996, in *Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers*, p. 997-998.
320. Lomax, J.D., 1996, DOE field test produces more oil, royalties from the Green River Formation: *Petroleum Engineer International*, v. 69, no. 8, p. 63-66,69.
321. Lomax, J., and Howard, A., 1994, New logging tool identifies permeability in shaley sands: *Oil and Gas Journal*, v. 92, no. 51, December 19, p. 104-108.
322. Looyestijn, W.J., 1994, *Le Log: The Log Analyst*, v. 36, no. 6, p. 6-7.
323. Looyestijn, W., 1996, Determination of oil saturation from diffusion NMR logs, paper SS, in *37th annual logging symposium transactions: Society of Professional Well Log Analysts*, 14 p.
324. Looyestijn, W., Schiet, M., and Runia, J., 1994, Successful field test of nuclear magnetic resonance (NMR) logging in the Rotterdam field, paper X, in *16th European formation evaluation symposium transactions: Society of Professional Well Log Analysts, Aberdeen Chapter*, 11 p.
325. Loren, J.D., 1972, Permeability estimates from NMR measurements: *Journal of Petroleum Technology*, v. 24, no. 8, p. 923-928.
326. Loren, J.D., and Robinson, J.D., 1969, Relations between pore size fluid and matrix properties and NMR measurements, SPE-2529: *Society of Petroleum Engineers, 44th Annual Fall Meeting Preprints*. Later published in 1970: *Society of Petroleum Engineers Journal*, v. 10, p. 268-278.
327. Lucas, A.J., Gibbs, S.J., Jones, E.W.G., Peyron, M., Derbyshire, J.A., and Hall, L.D., 1993, Diffusion imaging in the presence of static magnetic field gradients: *Journal of Magnetic*

- Resonance, v. 104, p. 273-282.
328. Lucas, A.J., Gibbs, S.J., Peyron, M., Pierens, G.K., Hall, L.D., Stewart, R.C., and Phelps, D.W., 1994, Pore geometry information via pulsed field gradient NMR: Magnetic Resonance Imaging, v. 12, no. 2, p. 249-251.
 329. Lucas, A.J., Peyron, M., Pierens, G.K., Carpenter, T.A., Hall, L.D., Stewart, R.C., Potter, G.F., and Phelps, D.W., 1993, A rigorous evaluation of the experimental requirements for performing quantitative porosity measurements by NMR, SPE-27420: Society of Petroleum Engineers, unsolicited manuscript, 59 p.
 330. Lucas, A.J., Pierens, G.K., Peyron, M., Carpenter, T.A., Hall, L.D., Stewart, R.C., Phelps, D.W., and Potter, G.F., 1993, Quantitative porosity mapping of reservoir rock cores by physically slice selected NMR, in Worthington, P.F., and Chardaire-Riviere, C., eds., Eurocas III--Advances in core evaluation management: Gordon and Breach Science Publishers, Philadelphia, p. 3-24.
 331. Lyne, A., Varini, G., and Ghilardotti, G., 1996, Determination of petrophysical properties of carbonate rocks by NMR relaxometry, SPE-36852, in SPE European petroleum conference proceedings, v. 1: Society of Petroleum Engineers, p. 331-339.
 332. Maddinelli, G., and Brancolini, A., 1996, MRI as a tool for the study of waterflooding processes in heterogeneous cores: Magnetic Resonance Imaging, v. 14, no. 7/8, p. 915-917.
 333. Maddinelli, G., Carati, C., Brancolini, A., Rossi, E., and Causin, E., 1993, NMR imaging of gelation processes inside rock cores, SPE-25219, in International symposium on oilfield chemistry proceedings: Society of Petroleum Engineers, p. 641-650.
 334. Maddinelli, G., Williams, J.L.A., and Taylor, D.G., 1991, A prospective model system to mimic the average NMR properties of water in a rock matrix: Magnetic Resonance Imaging, v. 9, no. 5, p. 743-746.
 335. Mahmood, S., Doughty, D. Tomutsa, L., and Honarpour, M.M., Pore level fluid imaging using high resolution nuclear magnetic resonance imaging and thin slice micromodels, paper SCA-9024, in Annual technical conference preprints: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, v. 3, v. 2, 20 p.
 336. Majors, P.D., Giveler, R.C., and Fukushima, E., 1989, Velocity and concentration measurements in multiphase flows by NMR: Journal of Magnetic Resonance, v. 85, p. 235-243.
 337. Majors, P.D., Li, P., and Peters, E.J., 1995, NMR imaging of immiscible displacements in porous media, SPE-30557, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology, part 1: Society of Petroleum Engineers, p. 263-278. Later published in 1997: SPE Formation Evaluation, v. 12, no. 3, p. 164-169.
 338. Majors, P.D., Smith, D.M., and Davis, P.J., 1991, Effect diffusivity measurement in porous media by NMR and radial imaging: Chemical engineering Science, v. 46, p. 3037-3043.
 339. Majors, P.D., Smith, J.L., Kovarik, F.S., and Fukushima, E., 1990, NMR spectroscopic imaging of oil displacement in dolomite: Journal of Magnetic Resonance, v. 89, p. 470-478.
 340. Mandava, S.S., Watson, A.T., and Edwards, C.M., 1990, NMR imaging of saturation during immiscible displacements: AIChE [American Institute of Chemical Engineering] Journal, v. 36, no. 11, p. 1680-1686.
 341. Mansfield, P., 1976, Proton spin imaging by nuclear magnetic resonance: Contemporary Physics, v. 17, p. 553-576.
 342. Mansfield, P., 1988, Imaging by nuclear magnetic resonance: Journal of Physics, Series E, v. 21, p. 18-30.
 343. Mansfield, P., Bowtell, R., Blackband, S., and Guilfoyle, D.N., 1992, Magnetic resonance imaging--applications of novel methods in studies of porous media: Magnetic Resonance Imaging, v. 10, p. 741-746.
 344. Mansfield, P., and Issa, B., 1994, Studies of fluid transport in porous rocks by echo-planar MRI: Magnetic Resonance Imaging, v. 12, no. 2, p. 275-278.
 345. Mansfield, P., and Issa, B., 1996, Fluid transport in porous rocks, part 1, EPI studies and a stochastic model of flow: Journal of Magnetic Resonance, series A, v. 122, p. 137-148.

346. Mansfield, P., and Issa, B., 1996, Fluid transport in porous rocks, part 2, Hydrodynamic model of flow and intervoxel coupling: *Journal of Magnetic Resonance*, series A, v. 122, p. 19-146.
347. Mardon, D., 1995, Basic physics of the nuclear magnetic resonance (NMR) logging tools, chapter 2, in Georgi, D.T., ed., *Nuclear magnetic resonance logging short course notes*, 36th annual symposium, Paris, France: Society of Professional Well Log Analysts, variously paginated.
348. Mardon, D., Gardner, J., Coates, G., and Vinegar, H., 1996, Experimental study of diffusion and relaxation of oil-water mixtures in model porous media, paper K, in 37th annual logging symposium transactions: Society of Professional Well Log Analysts, 14 p.
349. Mardon, D., Miller, D., Howard, A., Coates, G., Jackson, J., Spaeth, R., and Nankervis, J., 1996, Characterization of light hydrocarbon-bearing reservoirs by gradient NMR well logging--a Gulf of Mexico case study, SPE-36520, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 295-304. Also published in 1996 in condensed form: *Journal of Petroleum technology*, v. 48, no. 11, p. 1035-1036.
350. Mardon, D., Miller, D., Howard, A., Coates, G., Jackson, J., Spaeth, R., and Nankervis, J., 1996, Characterization of light hydrocarbon-bearing reservoirs by gradient NMR well logging--a Gulf of Mexico case study, SPE-36520, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 295-304.
351. Mardon, D., Prammer, M.G., and Coates, G.R., 1996, Characterization of light hydrocarbon reservoirs by gradient-NMR well logging: *Magnetic Resonance Imaging*, v. 14, nos. 7/8, p. 769-777.
352. Mardon, D., Prammer, M.G., Taicher, Z., Chandler, R.N. and Coates, G.R., 1995, Improved environmental corrections for MRIL pulsed NMR logs run in high-salinity boreholes, paper DD, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p.
353. Marschall, D.M., 1997, Laboratory MRI investigation in the effects of invert oil muds on primary MRI log determinations, SPE-38739, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology, part 2: Society of Petroleum Engineers, p. 203-215.
354. Marschall, D., 1997, Magnetic resonance technology and its applications in the oil and gas industry, part 2: *Petroleum Engineer International*, 70, no. 4, p. 65-70.
355. Marschall, D.M., and Coates, G., 1997, Laboratory MRI investigation in the effects of invert oil muds on primary MRI log determinations, paper SCA-9701, in 1997 international symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 11 p.
356. Marschall, D., Gardner, J., and Curby, F.M., 1997, MR laboratory measurements--requirements to assure successful measurements that will enhance MRI log interpretation, paper SCA-9704, in 1997 international symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 12 p.
357. Marschall, D., Gardner, J.S., Mardon, D., and Coates, G.R., 1995, Method for correlating NMR relaxometry and mercury injection data, paper SCA-9511, in International SCA symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 12 p.
358. Mattiello, D., Balzarini, M., Ferraccioli, L, and Brancolini, A., 1997, Calculation of constituent porosity in a dual-porosity matrix-- MRI and image analysis integration, paper SCA-9706, in 1997 international symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 10 p.
359. Maxwell, S., and Segel, S.L., 1985, A superfast NMR determination of the proton content of rocks, BHG5.6, in Expanded abstracts with biographies, 1985 technical program: Society of Exploration Geophysicists, p. 100-102. Later published in 1986: *Geophysics*, v. 51, no. 2, p. 438.
360. McKague, H.L., Hearst, J.R., Ward, R.L., and Burkhard, N. R., 1992, Nuclear magnetic resonance determination of the nonpore water content of zeolitic tuffs and its application to correction of

- epithermal-neutron-log-derived water content: *Nuclear Geophysics*, v. 6, no. 3, p. 359-366.
361. Meazza, O., Martera, M.D., Lyne, A., Taylor, M., Saunders, M.R., 1996, Porosity from cuttings--options and answers, paper SCA-9606, in International SCA symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 14 p.
362. Mendelson, K.S., 1986, Nuclear magnetic relaxation in fractal pores: *Physical Review, Series B*, v. 34, no. 9, p. 6503-6505.
363. Mendelson, K.S., 1986, Nuclear magnetic relaxation in porous media: *Journal of the Electrochemical Society*, v. 133, no. 3, p. 631-633.
364. Mendelson, K.S., 1989, Diffusion and relaxation in porous media: *Physica, Series A*, v. 157, p. 489-492.
365. Mendelson, K.S., 1990, Percolation model of nuclear magnetic relaxation in porous media: *Physical Review, Series B*, v. 41, p. 562-567.
366. Mendelson, K.S., 1991, Magnetic relaxation in porous media: *Magnetic Resonance Imaging*, v. 9, no. 5, p. 651-655.
367. Mendelson, K.S., 1991, Relation of spin-lattice relaxation; time to pore geometry and permeability: *Magnetic Resonance Imaging*, v. 9, p. 877-878.
368. Merrill, M.R., 1993, Local velocity and porosity measurements inside Casper Sandstone using MRI, paper 63E, in Spring national meeting proceedings: American Institute of Chemical Engineers, 15 p.
369. Merrill, M.R., and Jin, Z., 1994, Velocity measurements in natural porous media: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 345-348.
370. Meyer, G.H., Snowden, B.S., Jr., and Woessner, D.E., 1970, A tetrahedral model for pulsed nuclear magnetic resonance transverse relaxation application to the clay-water system: *Journal of Colloid and Interface Science*, v. 34, no. 1, p. 43-52.
371. Miknis, F.P., Smith, J.W., Maughan, E.K., and Maciel, G.E., 1982, Nuclear magnetic resonance--technique for direct nondestructive evaluation of source-rock potential: *AAPG Bulletin*, v. 66, no. 9, p. 1396-1401.
372. Miller, M.N., Paltiel, Z., Gillen, M.E., Granot, J., and Boouton, J.C., 1990, Spin-echo magnetic-resonance logging--porosity and free-fluid index, SPE-20561, in SPE annual technical conference exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 321-334.
373. Mirotnik, K., Allsopp, K., and Kantzas, A., 1997, Combination of NMR and ultracentrifuge techniques for effective carbonate reservoir characterization, paper SCA-9703, in 1997 international symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 10 p.
374. Mitra, P.P., Latour, L.L., Kleinberg, R.L., and Sotak, C.H., 1995, Pulsed field gradient NMR measurements of restricted diffusion and the return-to-the-origin probability: *Journal of Magnetic Resonance, Series A*, v. 114, p. 47.
375. Mitra, P.P., and Sen, P.N., 1992, Effects of microgeometry and surface relaxation on NMR pulsed-field-gradient experiments--simple pore geometries: *Physical Review, Series B*, v. 45, p. 143-156.
376. Mitra, P.P., and Sen, P.N., 1992, Effects of surface relaxation on NMR pulsed field gradient experiments in porous media: *Physica, Series A*, v. 186, p. 109-114.
377. Mitra, P.P., Sen, P.N., and Schwartz, L.M., 1993, Short-time behavior of the diffusion coefficient as a geometrical probe of porous media: *Physical Review, Series B*, v. 47, p. 8565-8574.
378. Mitra, P.P., Sen, P.N., Schwartz, L.M., and Le Doussal, P., 1992, Diffusion propagator as a probe of the structure of porous media: *Physical Review Letters*, v. 65, p. 3555.
379. Moore, M.A., and Akkurt, R., 1996, Nuclear magnetic resonance applied to gas detection in a highly laminated Gulf of Mexico turbidite invaded with synthetic oil filtrate, SPE-36521, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 305-310.

380. Morriss, C., 1995, CMR and basic log responses, chapter 7, in Georgi, D.T., ed., Nuclear magnetic resonance logging short course notes, 36th annual symposium, Paris, France: Society of Professional Well Log Analysts, variously paginated.
381. Morriss, C.E., Deutch, P., Freedman, R., McKeon, D., and Kleinberg, R.L., 1996, Operating guide for the Combinable Magnetic Resonance tool: *The Log Analyst*, v. 37, no. 6, p. 53-60.
382. Morriss, C.E., Freedman, R., Straley, C., Johnston, M., Vinegar, H.J., and Tutunjian, P.N., 1994, Hydrocarbon saturation and viscosity estimation from NMR logging in the Belridge diatomite, paper C, in 35th annual logging symposium transactions: Society of Professional Well Log Analysts, 24 p. Later published in 1997: *The Log Analyst*, v. 38, no. 2, p. 44-59.
383. Morriss, C.E., and Laverdiere, L., 1988, Nuclear magnetism log interpretation in shaly sand formations, paper EE, in 29th annual logging symposium transactions: Society of Professional Well Log Analysts, 22 p.
384. Morris, C.E., MacInnis, J., Freedman, R. Smaardyk, J., Straley, C., Kenyon, W.E., Vinegar, H.J., and Tutunjian, P.N., 1993, Field test of an experimental pulsed nuclear magnetism tool, paper GGG, in 34th annual logging symposium transactions: Society of Professional Well Log Analysts, 23 p.
385. Munn, K., and Smith, D.M., 1987, A NMR technique for the analysis of pore structure; numerical inversion of relaxation measurements: *Journal of Colloid and Interface Science*, v. 199, no. 1, p. 177-126.
386. Murphy, D.P., 1995, NMR logging and core analysis--simplified: *World Oil*, v. 216, no. 4, p. 65-70.
387. Murray, C.J., 1996, The use of nuclear magnetic resonance T2 relaxation measurements in the characterization of the porous microstructure and reservoir properties of sandstones: University of South Carolina, unpublished Ph.D. dissertation, 171 p.
388. Myal, F R, 1979, Residual oil saturation technology test in a consolidated sandstone reservoir, Bell Creek field, Powder River County, Montana, in 5th annual enhanced oil and gas proceedings, v. 2: U.S. Department of Energy, Report CONF-790805-P2, p. F-2/1--F-2/15.
389. Neretin, V.D., Belorai, Y.L., and Kononenko, I.Y., 1992, A study of permeability and porosity using nuclear magnetic resonance methods: *Nuclear Geophysics*, v. 6, no. 1, p. 121-124.
390. Neretin, V.D., and Chukhovichov, V.D., 1995, Interpretation of nuclear magnetic resonance data, in 36th annual logging symposium, Russian log interpretation workshop: Society of Professional Well Log Analysts, preprint, 21 p.
391. Nes, O.-M., and Skjetne, T., 1993, Shale porosities as determined by NMR, BG3.7, in 63rd annual meeting and international exhibition technical program expanded abstracts with biographies: Society of Exploration Geophysicists, p. 88-91.
392. Neuman, C.H., 1983, Application of the nuclear magnetism log and electromagnetic propagation tool to define steam potential in heavy oil reservoirs, in 12th annual convention proceedings, volume 2: Indonesian Petroleum Association, Jakarta, p. 265-273.
393. Neuman, C.H., and Brown, R.J.S., 1981, Applications of nuclear magnetism logging to formation evaluation, SPE-10108: Society of Petroleum Engineers, 56th annual meeting [San Antonio] preprint, 10 p. Later published in 1982: *Journal of Petroleum Technology*, v. 34, no. 12, p. 2853-2862.
394. Newbitt, G.J., de Groot, A., Fens, T.W., and Bonnie, J.H.M., 1991, Towards validation of porous media models using NMR imaging and image-analysis techniques: *Magnetic Resonance Imaging*, v. 9, p. 779-787.
395. Nigh, E.L., and Taylor, M., 1985, P-K--wellsite determination of porosity and permeability using drilled cutting, paper Z, in 10th formation evaluation symposium transactions: Canadian Well Logging Society, 16 p.
396. Nigh, E.L., Whittaker, A.H., and MacPherson, J.D., 1985, Statistical analysis of correlation of porosity and permeability determinations from well cuttings using a portable proton NMR apparatus with conventional core analysis and wireline log readings: *AAPG Bulletin*, v. 69, no. 4,

- p. 673.
397. Nikias, P.A., Jr., and Eyraud, L.E., 1962, Field examples of nuclear magnetism logging in the Poso Creek, Kern Front and Kern River fields of the San Joaquin Valley, SPE-434: Society of Petroleum Engineers 37th Annual Fall Meeting Preprints, 7 p.
 398. Norgaard, J.V., Olsen, D., Springer, N., and Reffstrup, J., 1995, Capillary pressure curves for low permeability chalk obtained by NMR imaging of core saturation profiles, SPE-30605, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 807-816.
 399. Nurmi, R., and Standen, E., 1997, Carbonates, the inside story: Middle East Well Evaluation Review, no. 18, p. 27-41.
 400. Ohen, H.A., Ajufo, A., and Curby, F.M., 1995, A hydraulic (flow) unit based model for the determination of petrophysical properties from NMR relaxation measurements, SCA-9514, in International SCA symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 12 p. Also published in 1995 as, SPE-30626, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 983-996.
 401. Ohen, H.A., Ajufo, A.P., and Enwere, P.M., 1996, Laboratory NMR relaxation measurements for the acquisition of calibration data for NMR logging tools, SPE-35683, in 66th SPE western region meeting proceedings: Society of Petroleum Engineers, p. 329-342.
 402. Olesen, J.-R., Liu, C., Zeng, W., Mao, K., and Zhang, Z., 1997, Evaluation of viscous oil-bearing, water flooded reservoirs, paper T, in 3rd well logging symposium of Japan proceedings: Society of Professional Well Log Analysts, Japan Chapter, 14 p.
 403. Olsen, D., 1994, MRI of a waterflood on a reservoir chalk sample: Magnetic Resonance Imaging, v. 12, no. 2, p. 215-216.
 404. Olsen, D., Topp, S., Stensgaard, A., Norgaard, J.V., and Reffstrup, J., 1996, Quantitative 1D saturation profiles on chalk by NMR: Magnetic Resonance Imaging, v. 14, no. 7/8, p. 847-851.
 405. Oosting, P.H., and Trappeniers, N.J., 1971, Proton spin-lattice relaxation and self-diffusion in methanes, part 3--Interpretation of proton spin-relaxation experiments: Physica, v. 51, p. 395-417.
 406. Oosting, P.H., and Trappeniers, N.J., 1971, Proton spin-lattice relaxation and self diffusion in methanes, part 4--Self-diffusion in methanes: Physica, v. 51, p. 418-431.
 407. Oraby, M., Chafai, N., Hussing, R.B., Massengill, D.R., Clarker, J.S., and Pletcher, D., 1997, A new NMR interpretation technique using error minimization with variable T2 cutoff, SPE-38734, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology, part 2: Society of Petroleum Engineers, p. 175-181.
 408. Oraby, M.E., and Eubanks, D.L., 1997, Determination of irreducible water saturation using magnetic resonance imaging logs (MRIL)--a case study from east Texas, USA, SPE-37772, in 10th Middle East oil show proceedings, v. 2: Society of Petroleum Engineers, p. 155-162.
 409. Oren, P.E., Rueslatten, H.G., Skjetne, T., and Buller, A.T., 1994, Some advances in NMR characterization of reservoir sandstones, in Aasen, J.O., Berg, E., Buller, A.T., Hjelmeland, O., Holt, R.M., Kleppe, J., and Torsaeter, O., eds., North Sea oil and gas reservoirs III: Kluwer Academic Publishers, London, p. 307-316.
 410. Osment, P.A., Packer, K.J., Taylor, M.J., Attard, J.J., Carpenter, T.A., Hall, L.D., Nerrod, N.J., and Doran, S.J., 1991, NMR imaging of fluids in porous solids: Philosophical Transactions of the Royal Society of London, Series A, v. 333, no. 1632, p. 441-452.
 411. Packer, K.J., and Zelaya, F.O., 1989, Observations of diffusion of fluids in porous media by pulsed field gradient NMR: Colloids Surfaces, v. 36, p. 221-227.
 412. Patz, S., Stromski, M., Hrovat, M., Straley, C., Schwartz, L., 1991, Application of single species chemical shift imaging to sandstone core: Magnetic Resonance Imaging, v. 9, p. 797-802.
 413. Pelletier, R., 1987, NMR applications in the Belly River formation of southern Alberta, paper W, in 11th formation evaluation symposium transactions: Canadian Well Logging Society, 14 p.
 414. Peters, A.M., Robyr, P.S., Bowtell, R.W., and Mansfield, P., 1996, Echo-planar microscopy of

- porous rocks: *Magnetic Resonance Imaging*, v. 14, no. 7/8, p. 875-877.
415. Petricola, M.J.C., and Takezaki, H., 1996, Nuclear magnetic resonance logging--can it minimize well testing?, SPE-36328, in 7th Abu Dhabi international petroleum conference proceedings: Society of Petroleum engineers, p. 740-748.
416. Petricola, M.J.C., and Watfa, M., 1997, Porosity in carbonates--how wet is too wet?: *Petroleum Engineer International*, v. 70, no. 3, p. 23-27.
417. Peyron, M., Pierens, G.K., Lucas, A.J., Hall, L.D., Potter, G.F., Stewart, R.C., and Phelps, D.W., 1994, Strategies for overcoming linewidth limitations in quantitative petrophysical NMR measurements: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 295-298.
418. Phen, H.A., Ajufo, A., and Curby, F.M., 1995, A hydraulic (flow) unit based model for the determination of petrophysical properties from NMR relaxation measurements, paper SCA-9514, in International SCA symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 12 p.
419. Potter, Kleinberg, R.L., Brockman, and MacFarlane, 1996, Assay for bacteria in porous media by diffusion weighted NMR: *Journal of Magnetic Resonance, Series B*, v. 113, p. 9.
420. Prammer, M.G., 1994, NMR pore size distributions and permeability at the well site, SPE-28368, in SPE annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 55-64.
421. Prammer, M., 1995, Principles of signal processing--NMR data and T2 distributions, chapter 4, in Georgi, D.T., ed., Nuclear magnetic resonance logging short course notes, 36th annual symposium, Paris, France: Society of Professional Well Log Analysts, variously paginated.
422. Prammer, M.G., Drack, E.D., Bouton, J.C., Gardner, J.S., Coates, G.R., Chandler, R.N., and Miller, M.N., 1996, Measurements of clay-bound water and total porosity by magnetic resonance logging, SPE-36522, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 111-118. Also published in 1996: *The Log Analyst*, v. 37, no. 5, p. 61-69.
423. Prammer, M.G., Mardon, D., Coates, G.R., and Miller, M.N., 1995, Lithology-independent gas detection by gradient-NMR logging, SPE-30562, in Annual technical conference and exhibition proceedings, v. omega, Formation evaluation and reservoir geology: Society of Petroleum Engineers, p. 325-336.
424. Praxmeyer, T., 1967, Range in nuclear magnetic logging: *Acta Geophysica Polonica*, v. 15, no. 2, p. 105-109.
425. Praxmeyer, T., 1967, Reception of the nuclear magnetism logging signal: *Acta Geophysica Polonica*, v. 15, no. 3, p. 217-224.
426. Prieto, M., Barbato, R., Sinha, A., and Gomez, G., 1997, Conciliation of core data and log derived permeability from nuclear magnetic resonance and Stoneley wave in northern Monagas area, eastern Venezuela, paper WW, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 14 p.
427. Puckett, D.A., and Sims, R.J., 1986, NMR imaging as an aid to experimental enhanced oil recovery: United Kingdom Atomic Energy Agency, Atomic Energy Establishment Winfrith Report No. AEEW-R-2103, 36 p.
428. Relling, O., Eidesmo, T., Flolo, L.H., and Rueslatten, H., 1996, NMR logging in a North Sea gas well, paper GGG, in 37th annual logging symposium transactions: Society of Professional Well Log Analysts, 10 p.
429. Riepe, L., Schopper, J.R., and Wehr, R., 1986, Permeability estimations from nuclear measurements (NMR-, spectral GR-data), paper X, in 10th Europe formation evaluation symposium transactions: Society of Professional Well Log Analysts, Aberdeen Chapter, 15 p.
430. Robinson, J.D., Loren, J.D., Vajnar, E.A., and Hartman, D.E., 1972, Determining residual oil with the nuclear magnetism log, SPE-3797, in Symposium on improved oil recovery, proceedings: Society of Petroleum Engineers. Later published in 1974: *Journal of Petroleum Technology*, v. 26, no. 2, p. 226-236.

431. Robinson, J.D., and Senturia, S.D., 1970, Nuclear spin-lattice relaxation of liquids confined in porous solids, SPE-2870: Society of Petroleum Engineers Journal, v. 10, no. 3, p. 237-244.
432. Robinson, M.A., 1991, Magnetic resonance imaging for the measurement of oil core porosity distributions, SPE-23582: Society of Petroleum Engineers, unsolicited paper, 31 p.
433. Robinson, M.A., Deans, H.A., and Bansal, S., 1992, Determination of oil core flow velocities and porosities using MRI, SPE-23960, in SPE Permian Basin oil and gas recovery conference, proceedings: Society of Petroleum Engineers, p. 261-268.
434. Robinson, M.A., and Edelstein, W.A., 1991, Fluid velocities in oil cores during water injection, SCA-9113, in 5th annual technical conference Preprints, v. 3: Society of Professional Well Log Analysts, Society of Core Analysts chapter-at-Large, 13 p.
435. Roestenburg, J.W., 1996, The combined application and utilisation of CMR and FMI wireline technology in Wonnich-1: APPEA [Australasian Petroleum Production Exploration Association] Journal, v. 36, part 1, p. 202-208.
436. Rothwell, W.P., and Vinegar, H.J., 1985, Petrophysical applications of NMR imaging: Applied Optics, v. 24, no. 23, p. 3969-3972.
437. Rueslatten, H., Eidesmo, T., Lehne, K.A., and Relling, O.M., 1996, The use of NMR spectroscopy to validate NMR logs from deeply buried reservoir sandstones, paper SCA-9621, in International SCA symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 11 p.
438. Rueslatten, H., Oren, P.-E., Robin, M., Rosenberg, E., and Cuiec, L., 1994, A combined use of cryo-SEM and NMR-spectroscopy for studying the distribution of oil and brine in sandstones, SPE-27804, in SPE/DOE 9th symposium on improved oil recovery proceedings: Society of Petroleum Engineers, p. 243-254.
439. Sakurai, S. and Loucks, R.G., and Gardner, J.S., 1995, NMR core analysis of Lower San Andres/Glorieta/Upper Clear Fork (Permian) carbonates--central basin platform, west Texas, paper M, in 36th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p.
440. Sanders, J.K., and Huntess, A., 1987, Modern NMR spectroscopy: Oxford University Press.
441. Sandia National Laboratory, 1979, Logging techniques based on dispersive electrical parameters: Report No. SAND 79-0878C/HDM.
442. Sandor, R., Bonnie, R., Tutunjian, P., and Ferris, J., 1994, Frequency dependence of NMR measurements on rock samples, paper L, in 16th European formation evaluation symposium transactions: Society of Professional Well Log Analysts, Aberdeen Chapter, 8 p.
443. Sandor, R.K.J. and Looyestijn, W.J., 1995, NMR logging--the new measurement: Shell International Petroleum Maatschappij, The Hague, The Netherlands, 8 p.
444. Santarelli, F.J., Marsala, A.F., Brignoli, M., Rossi, E. and Bona, N., 1996, Formation evaluation from logging on cuttings, SPE-36851, in European petroleum conference [EUROPEC] proceedings, v. 1: Society of Petroleum Engineers, p. 319-329.
445. Saraf, D.N., and Fatt, I., 1967, Three-phase relative permeability measurements using a nuclear magnetic resonance technique for estimating fluid saturation, SPE-1760: Society of Petroleum Engineers Journal, v. 7, no. 3, p. 235-242.
446. Schlumberger, 1997, CMR Combinable Magnetic Resonance Tool--User's Guide: Schlumberger Wireline and Testing, Houston, Texas, Document No. SMP-7059, variously paginated.
447. Schmidt, E.J., 1987, Nuclear magnetic resonance and the broadband acoustic response of porous rocks: Stanford University, Stanford, California, unpublished Ph.D. dissertation, 198 p.
448. Schmidt, E.J., Velasco, K., and Nur, A.M., 1985, Quantifying solid-fluid interfacial phenomena in porous rock with proton nuclear magnetic resonance: Proceedings of the Electrochemistry Society, v. 85-88, p. 292-304. Later published in 1986: Journal of Applied Physics, v. 59, no. 8, p. 2788-2797.
449. Schwartz, L.M., Bergman, D.J., Dunn, K.J., and Mitra, P.P., 1996, Self-diffusion in periodic porous media--a comparison of numerical simulation and eigenvalue methods: Magnetic

- Resonance Imaging, v. 14, no. 7/8, p. 737-743.
450. Schwartz, L.M., Sen, P.B., and Mitra, P.P., 1994, Simulations of pulsed field gradient spin-echo measurements in porous media: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 241-244.
 451. Schwartz, L.M., Wilkinson, D.J., Kostek, S., Johnson, D.L., and Banavar, J.R., 1991, Nuclear magnetism and transport in porous media: *Magnetic Resonance Imaging*, v. 9, p. 657-662.
 452. Seevers, D.O., 1966, A nuclear magnetic method for determining the permeability of sandstone, paper L, in 7th annual logging symposium transactions: Society of Professional Well Log Analysts, 14 p.
 453. Seevers, D.O., 1972, A method of determining permeability by means of a pulsed NMR and an application of the method to the study of a Gulf Coast supernormally pressured well, SPE-3847, in, 3rd abnormal subsurface pore pressure symposium preprints: Society of Petroleum Engineers, p. 67-74.
 454. Sen, P.N., Schwartz, L.M., and Mitra, P.P., 1994, Probing the structure of porous media using NMR spin echoes: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 227-230.
 455. Sen, P.N., Schwartz, L.M., and Mitra, P.P., and Halperin, B.I., 1994, Surface relaxation and the long-time diffusion coefficient in porous media--periodic geometries: *Physical Review, Series B*, v. 49, p. 215.
 456. Sen, P.N., Straley, C., Kenyon, W.E., and Whittingham, M.S., 1990, Surface-to-volume ratio, charge density, nuclear magnetic relaxation, and permeability in clay-bearing sandstones: *Geophysics*, v. 55, no. 1, p. 61-69.
 457. Senturia, S.D., and Robinson, J.D., 1970, Nuclear spin-lattice relaxation of liquids confined in porous solids: *Society of Petroleum Engineers Journal*, v. 10, no. 3, p. 237-244.
 458. Sercombe, W.J., and Anderson, B.R., 1990, Net-pay application from nuclear magnetic resonance, Nugget Sandstone, Wyoming thrust belt, SCA-9026, in Annual technical conference preprints: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, v. 3, v. 2, 22 p.
 459. Setser, G., and Williams, M.R., 1985, Measurement of remaining oil saturation in northern Michigan using nuclear magnetism log data and pressure core, SPE-14276: Society of Petroleum Engineers, 60th annual meeting preprint, 10 p.
 460. Sezginer, A., Griffin, D.D., Kleinberg, R.L., Fukuhara, M., and Dudley, D.G., 1993, RF sensor of a novel NMR apparatus: *Journal of Electromagnetic Waves and Applications*, v. 7, no. 1, p. 13-30.
 461. Sezginer, A., Kleinberg, R.L., Fukuhara, M., and Latour, L.L., 1991, Very rapid simultaneous measurement of nuclear magnetic resonance spin-lattice relaxation time and spin-spin relaxation time: *Journal of Magnetic Resonance*, v. 92, no. 3, p. 504-527.
 462. Sharma, R.K., Das, T.K., Sood, S.K., and Kotho, J.M., 1994, Paramagnetism resonance--a tool for logging, OSEA-94109, in 10th offshore South East Asia conference and exhibition preprints: American Association of Petroleum Geologists, p. 849-855.
 463. Shushakov, O.A., 1996, Groundwater NMR in conductive water: *Geophysics*, v. 61, no. 4, p. 998-1006.
 464. Singer, J., Johnston, L., Kleinberg, R., and Flaum, C., 1997, Fast NMR logging for bound fluid and permeability, paper YY, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 13 p.
 465. Skjetne, T, Southon, T.E., Hafskjold, B., Selle, O., Svorstol, I., Buller, A.T., Rueslatten, H., Brayshaw, A., and Kalam, M.Z., 1991, Nuclear magnetic resonance studies of reservoir core plugs--preliminary investigation of the influence of mineralogy on T1: *Magnetic Resonance Imaging*, v. 9, p. 673-679.
 466. Skopec, R.A., Foot, J., and Marion, R., 1996, Petrophysical innovations and their influence on exploration and production in NW Europe, in Glennie, K., and Hurst, A., eds., AD1995--NW Europe's hydrocarbon industry: Geological Society [London], p. 177-184.
 467. Slichter, C.P., 1990, Principles of magnetic resonance: Springer-Verlag, New York.

468. Smith, D.M., Holt, T.E., Gallegos, D.P., and Stermer, D.L., 1987, Pore structure analysis via NMR, mercury porosimetry a, and dynamic methods: *Advances in Ceramic Powder Science*, v. 21, p. 779-791.
469. Smith, S., and Pickens, T., 1997, Texaco looks at magnetic resonance logs vs. coring in GOM: *World Oil*, v. 218, no. 11, p. 114-116.
470. Solanet, F., Khatchikian, A., and Breda, E., 1997, NMR as standard porosity tool in the San Jorge Basin--log responses and applications, SPE-38735, in *Annual technical conference and exhibition proceedings*, v. omega, Formation evaluation and reservoir geology, part 2: Society of Petroleum Engineers, p. 183-195.
471. Sorenson, R.P., 1992, Recognition and evaluation of pressure-depleted reservoirs, paper LL, in *33rd annual logging symposium transactions: Society of Professional Well Log Analysts*, 14 p. Later published in 1994: *Houston Geological Society Bulletin*, v. 37, no. 2, October, p. 17-21.
472. Spinler, E.A., 1996, Methodology for determining the distribution of wettability from laboratory and well log measurements for a microcrystalline/microfossiliferous carbonate field, SCA-9618, in *International SCA symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large*, 10 p.
473. Stallmach, F., Appel, M., and Thomann, H., 1996, Irreducible fluid saturation determined by pulsed field gradient NMR, paper SCA-9620, in *International SCA symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large*, 15 p.
474. Stewart, W.F., 1996, *Le Log: The Log Analyst*, v. 37, no. 3, p. 12-13.
475. Stonard, S.W., LaTorraca, G.A., and Dunn, K-J., 1996, Effects of magnetic susceptibility contrasts on oil and water saturation determinations by NMR T2 laboratory and well log measurements, paper SCA-9619, in *International SCA symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large*, 12 p.
476. Stone, W.E.E., 1982, The use of NMR (nuclear magnetic resonance), in *The study of clay minerals, advanced techniques for clay mineral analysis: Elsevier Scientific Publishing Company, New York, Developments in Sedimentology Series No. 34*, p. 77-112.
477. Straley, C., 1991, New horizons from the lab to the field: *Magnetic Resonance Imaging*, v. 9, p. 875-876.
478. Straley, C., 1997, An experimental investigation of methane in rock materials, paper AA, in *38th annual logging symposium transactions: Society of Professional Well Log Analysts*, 14 p.
479. Straley, C., Matteson, A., Feng, S., Schwartz, L., Kenyon, W.E., and Banavar, J.R., 1987, Magnetic resonance, digital image analysis, and permeability of porous media: *Applied Physics Letters*, v. 51, no. 15, p. 1146-1148.
480. Straley, C., Morriss, C.E., Kenyon, W.E., and Howard, J.J., 1991, NMR in partially saturated rocks: laboratory insights on free fluid index and comparison with borehole logs, paper CC, in *32nd annual logging symposium transactions: Society of Professional Well Log Analysts*, 25 p. Later published in 1994: *The Log Analyst*, v. 36, no. 1, p. 40-56.
481. Straley, C., Rossini, D., Schwartz, L.M., Stromski, M.E., Hrovat, M., and Patz, S., 1992, Particle filtration in sandstone cores--application of a chemical shift imaging technique, SCA-9217, in *6th annual technical conference preprints [33rd SPWLA annual logging symposium transactions, v. 3]: Society of Core Analysts, Society of Professional Well Log Analysts Chapter-at-Large*, 24 p. Later published in 1995: *The Log Analyst*, v. 36, no. 2, p. 42-51.
482. Straley, C., Rossini, D., Schwartz, L.M., Stromski, M.E., Hrovat, M., and Patz, S., 1994, Chemical shift imaging of particle filtration in sand: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 313-315.
483. Straley, C., Rossini, D., Vinegar, H., Tutunjian, P., and Morriss, C., 1994, Core analysis by low field NMR, paper SCA-9404, in *SCA international symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts, Chapter-at-Large*, p. 43-56. Later published in 1997: *The Log Analyst*, v. 38, no.2, p. 84-93.
484. Straley, C., and Schwartz, L.M., 1996, Transverse relaxation in random bead packs--comparison

- of experimental data and numerical simulations: *Magnetic Resonance Imaging*, v. 14, no. 7/8, p. 999-1002.
485. Su, M.Y., Nalcioğlu, O., 1993, Spin-lattice and spin-spin relaxation in porous media--a generalized two site relaxation model: *Journal of Colloid and Interface Science*, v. 160, no. 2, p. 332-337.
486. Sully, J.P., and Campanella, R.G., 1995, Evaluation of in situ anisotropy from crosshole and downhole shear wave velocity measurements: *Geotechnique*, v. 45, no. 2, p. 267-282.
487. Sutyono, S., 1995, Magnetic resonance image log use in evaluation of low resistivity pay in the Attaka field, paper IPA95-2.2-056, in 24th annual convention proceedings, v. 2: Indonesian Petroleum Association, Jakarta, 13 p.
488. Szemeredy, P., 1986, Problem of direct task in nuclear magnetism logging: *Acta Geodetica Geophysica, and Montanistica Hungarica*, v. 24, no. 1-2, p. 127-141.
489. Taicher, Z., Coates, G., Gitartz, Y., and Berman, L., 1994, A comprehensive approach to studies of porous media (rocks) using a laboratory spectrometer and logging tool with similar operating characteristics: *Magnetic Resonance Imaging*, v. 12, no. 2, p. 285-289.
490. Tariq, S.M., Denoo, S., and Cordes, M., 1997, Permeability estimation from modern NMR logging measurements; experiences and applications in the Rocky Mountain area, SPE-38378, in SPE Rocky Mountain regional meeting/Geological Association field conference proceedings: Society of Petroleum Engineers.
491. Thompson, A.H., Katz, A.J., and Krohn, C.E., 1987, The microgeometry and transport properties of sedimentary rock: *Advances in Physics*, v. 36, no. 5, p. 625-694.
492. Thompson, A.H., Sinton, S.W., Huff, S.L., Katz, A.J., Raschke, R.A., and Gist, G.A., 1989, Deuterium magnetic resonance and permeability in porous media: *Journal of Applied Physics*, v. 65, p. 3259-3263.
493. Timur, A., 1964, Studies of multicomponent and multiphase fluid flow in porous media through use of a nuclear magnetic resonance method, SPE-1066: Society of Petroleum Engineers, unsolicited paper, 18 p..
494. Timur, A., 1967, Pulsed nuclear magnetic resonance studies of porosity, movable fluid and permeability of sandstones, SPE-2045: Society of Petroleum Engineers, 42nd Annual Meeting Preprint. Later published in 1969: *Journal of Petroleum Technology*, v. 21, no. 6, p. 775-786.
495. Timur, A., 1968, An investigation of permeability, porosity, and residual water saturation relationships for sandstone reservoirs, paper J, in 9th annual well logging symposium transactions: Society of Professional Well Log Analysts, 18 p. Later published in 1968: *The Log Analyst*, v. 9, no. 4, July-August, p. 8-17. Later reprinted in 1994, in *Petrophysics*: Society of Petroleum Engineers Reprint Series No. 39, p. 128-137.
496. Timur, A., 1968, Effective porosity and permeability of sandstones investigated through nuclear magnetic principles, paper K, in 9th annual logging symposium transactions: Society of Professional Well Log Analysts, 18 p. Later published in 1969: *The Log Analyst*, v. 10, no. 1, January-February, p. 3-11.
497. Timur, A., 1972, Nuclear magnetic resonance study of carbonates, paper N, in 13th annual logging symposium transactions: Society of Professional Well Log Analysts, 15 p. Later published in 1972: *The Log Analyst*, v. 13, no. 5, p. 3-11.
498. Timur, A., Hemphkins, W.B., and Massey, G.R., 1971, Analysis of sidewall samples by nuclear magnetic resonance methods, paper T in 12th annual logging symposium transactions: Society of Professional Well Log Analysts, 21 p. Also published in 1971: *The Log Analyst*, v. 12, no. 6, p. 13-19.
499. Trushkin, D.V., Shushakov, O.A., and Legchenko, A.V., 1995, Surface NMR applied to an electroconductive medium: *Geophysical Prospecting*, v. 43, no. 5, p. 623-633.
500. Tulbovich, B.I., Borsutskiy, Z.R., and Zlobin, A.A., 1984, Investigation of infiltration-capacity properties and structural characteristics of oil-bearing rocks by the NMR impulse method: *International Geological Review*, v. 26, no. 4, p. 456-461.

501. Tutunjian, P.N., and Vinegar, H.J., 1990, Petrophysical applications of automated proton high-resolution NMR spectroscopy, SCA-9020, in 4th annual technical conference preprints: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, v. 2 (1990), 18 p. Later published in 1992: *The Log Analysts*, v. 33, no. 2, p. 136-144.
502. Tutunjian, P.N., Vinegar, H.J., and Edelstein, W.A., 1991, Automated core analysis by ¹H NMR spectroscopy: *Magnetic Resonance Imaging*, v. 9, no. 5, p. 859-864.
503. Tutunjian, P.N., Vinegar, H.J., and Ferris, J.A., 1991, Nuclear magnetic resonance imaging of sodium-23 in cores, SCA-9111, in 5th annual technical conference preprints: Society of Core Analysts, Society of Professional Well Log Analysts Chapter-at-Large, v. 3, 18 p. Later published in 1993: *The Log Analyst*, v. 34, no. 3, p. 11-19.
504. Tutunjian, P.N., Vinegar, H.J., Ferris, J.A., and Nesbitt, G.J., 1993, Characterization of pore microgeometry by NMR diffusion measurements, paper II, in 34th annual logging symposium transactions: Society of Professional Well Log Analysts, 13 p.
505. Vinegar, H.J., 1986, X-ray CT and NMR imaging of rocks, SPE-15277: *Journal of Petroleum Technology*, v. 38, no. 3, p. 257-259. Later reprinted in 1994, in *Petrophysics: Society of Petroleum Engineers Reprint Series No. 39*, p. 115-117.
506. Vinegar, H., 1995, Relaxation mechanisms, chapter 3, in Georgi, D.T., ed., *Nuclear magnetic resonance logging short course notes*, 36th annual symposium, Paris, France: Society of Professional Well Log Analysts, variously paginated.
507. Vinegar, H., 1995, NMR log interpretation with examples, chapter 8, in Georgi, D.T., ed., *Nuclear magnetic resonance logging short course notes*, 36th annual symposium, Paris, France: Society of Professional Well Log Analysts, variously paginated.
508. Vinegar, H.J., Tutunjian, P.N., Crabtree, P.T., Raffaldi, F.J., DiFoggio, R., and Edelstein, W.A., 1989, NMR spectroscopy of tight, gypsum-bearing carbonates, paper HH, in 30th annual logging symposium transactions: Society of Professional Well Log Analysts, 15 p. Later published in 1991: *The Log Analyst*, v. 32, no. 5, p. 527-535.
509. Vinegar, H.J., Tutunjian, P.N., Edelstein, W.A., and Roemer, P.B., 1989, ¹³C NMR of whole cores, SPE-19590, in SPE annual technical conference and exhibition proceedings, v. omega, *Formation evaluation and reservoir geology: Society of Petroleum Engineers*, p. 201-212. Later published in 1991: *SPE Formation Evaluation*, v. 6, no. 2, p. 183-189.
510. Vinegar, H.J., Tutunjian, P.N., Edelstein, W.A., and Roemer, P.B., 1989, Determining carbonate content of cores by ¹³C NMR, SCA-8901, part III, paper A, in Annual technical conference reprints, v. 1 (1987-1989): Society of Professional Well Log Analysts, Society of Core Analysts SPWLA Chapter-at-Large, 18 p.
511. Vogeley, J.R., and Moses, C.O., 1992, ¹H NMR relaxation and rock permeability: *Geochimica, et Cosmochimica Acta*, v. 56, p. 2947-2953.
512. Walsgrove, T., Stromberg, S., Lowden, B., and Basan, P., 1997, Integration of nuclear magnetic core analysis and nuclear magnetic resonance logs--an example from the North Sea, paper UU, in 38th annual logging symposium transactions: Society of Professional Well Log Analysts, 12 p. A similar paper also published in 1997 as Basan, P., Lowden, B., and Stromberg, S., *Developing strategies using core analysis for calibrating the NMR log: DiaLog [London Petrophysical Society Newsletter]*, v. 5, no. 6, p. 4-6.
513. Wang, W., Land, D., and Liu, W., 1996, The application of NMR imaging to the studies of enhanced oil recovery in China: *Magnetic Resonance Imaging*, v. 14, no. 7/8, p. 951-953.
514. Wardlaw, N.C., 1983, Rock texture, pore structure wettability--neglected parameters in well log interpretations of fluid saturations?: *Canadian Well Logging Society Journal*, v. 12, p. 89-98. Later reprinted in 1994, in *Petrophysics: Society of Petroleum Engineers Reprint Series No. 39*, p. 105-114.
515. Warrilow, I.M., Ugueto-Cabera, G.A., Nawawi, H., Sibbit, A., 1997, Nuclear magnetic resonance--a useful complement to reservoir characterisation of Brunei deltaic sequences, SPE-38025, in *Asia Pacific oil and gas conference proceedings: Society of Petroleum Engineers*, p. 91-98.

516. White, J., and Bedford, J., 1997, Nuclear magnetic resonance applications in the North Sea: DiaLog [London Petrophysical Society Newsletter], v. 5, no. 6, p. 6-8.
517. Widjanarko, W., 1996, Integrating nuclear magnetic resonance logging data with traditional down hole petrophysical data to optimize new development wells strategies in the Bravo field offshore north west Java, paper IPA96-2-2-162, in 25th annual convention proceedings, v. 2: Indonesian Petroleum Association, Jakarta, p. 205-213.
518. Wilkenson, D.J., Johnson, D.L., and Schwartz, L.M., 1991, Nuclear magnetic relaxation in porous media; the role of mean lifetime: Physical Review, Series B, v. 44, p. 4960.
519. Williams, C.E., and Fung, B.M., 1982, Determination of wettability by hydrocarbons of small particles by deuteron T1 measurements: Journal of Magnetic Resonance, v. 50, p. 71-80.
520. Williams, J.L.A., 1992, The application of MRI to enhanced oil recovery studies: University of Surrey, England, unpublished Ph.D., dissertation.
521. Williams, J.L.A., Maddinelli, G., Taylor, D.G., Enwere, P., and Archer, J.S., 1991, Contribution of NMR imaging technique in the study of polyphasic flow in porous media: Magnetic Resonance Imaging, v. 9, no. 5, p. 869-874.
522. Williams, J.L.A., and Taylor, D.G., 1994, Measurements of viscosity and permeability of two phase miscible fluid flow in rock cores: Magnetic Resonance Imaging, v. 12, no. 2, p. 317-318.
523. Williams, J.L.A., Taylor, D.G., Maddinelli, G., Enwere, P., and Archer, J.S., 1991, Visualisation of fluid displacement in rock cores by NMR imaging: Magnetic Resonance Imaging, v. 9, p. 767-773.
524. Withjack, E.M., and Durham, J.R., 1995, Characterization and saturation determination of geothermal rock by CT and NMR, paper SCA-9515, in International SCA symposium proceedings: Society of Professional Well Log Analysts, Society of Core Analysts Chapter-at-Large, 10 p.
525. Wyman, R.E., 1962, Nuclear magnetism field test results, paper 12, in 3rd annual logging symposium transactions: Society of Professional Well Log Analysts, 14 p.
526. Wyman, R.E., 1962, The NML--what Shell is learning about it: Petroleum Management, v. 34, no. 12, p. 82-89.
527. Wyman, R.E., 1978, Logging methods, chapter 5, in Determination of residual oil saturation: Interstate Oil Compact Commission, Oklahoma City, p. 89-116. Later reprinted in 1991, in Canadian Heavy Oil Association reservoir handbook: Canadian Heavy Oil Association, Calgary, p. 69-81.
528. Xiao, L., Du, Y., Ye, C., 1994, ^{13}C Longitudinal relaxation of fluids saturated in oil reservoir cores by MAS (magic angle spinning) NMR: Journal of Colloid and Interface Science, v. 164, no. 2, p. 495-497.
529. Yousupov, M.Z., Chukhvichyov, V.D., Voronkov, L.N., Donovan, G.M., Koshlyak, V.A., 1979, Logging methods in studying oil beds flooded with fresh water, paper R, in 6th European formation evaluation symposium transactions: Society of Professional Well Log Analysts, London Chapter, 15 p.